

CSE 20 Lecture 15

Analysis: Inclusion- Exclusion Technique

CK Cheng

Analysis: Inclusion – Exclusion Principle

Ex: A bakery makes only $M=3$ kinds of cookies. Find the number of ways a person can buy $r=4$ of the cookies.

$x+y+z = 4$ (each variable represents the cookie in that flavor)

$$C(4+2, 2) = (6*5)/2! = 15$$

$$(C(n + m - 1, m - 1),$$

$n = \text{\#cookies},$

$m = \text{\#kinds of cookies})$

X	Y	Z
0	0	4
0	1	3
0	2	2
0	3	1
0	4	0
1	0	3
1	1	2
...

Cont. (from last slide)

Ex: Buy 2 cookies out of 3 flavors

$$x+y+z = 2$$

$$C(2+2, 2) = C(4,2)$$

$$= (4*3)/2! = 6$$

Id	X	Y	Z
0	0	0	2
1	0	1	1
2	0	2	0
3	1	0	1
4	1	1	0
5	2	0	0

Cont. (from last slide)

Ex: Buy 8 cookies out of 4 kinds of flavors.

$$x_1 + x_2 + x_3 + x_4 = 8$$

$$C(8+3, 3) = 11! / (8!3!) = (11 \cdot 10 \cdot 9) / 3! = 165.$$


Ex: Buy 20 cookies out of 5 kinds of flavors.

$$x_1 + x_2 + x_3 + x_4 + x_5 = 20$$

$$C(20+4, 4) = 24! / (20!4!) = (24 \cdot 23 \cdot 22 \cdot 21) / 4!$$

Ex: Possible number of Routing Paths

Assume that the streets are on a grid and the walking distance must be the shortest. Find the number of ways to walk from (0,0) to (6,3).



View from vertical direction:

$$Y_0 + Y_1 + Y_2 + Y_3 + Y_4 + Y_5 + Y_6 = 3$$
$$\rightarrow C(3+6, 6) = C(9, 6) = 9!/(6!3!)$$

View from horizontal direction:

$$X_0 + X_1 + X_2 + X_3 = 6$$
$$\rightarrow C(6+3, 3) = C(9, 6) = 9!/(6!3!)$$

Cont. (from last slide)

Ex: On a grid street, find #shortest paths from (1,3) to (2,5).

$$\text{Formulations: } x_1 + x_2 = (5-3)$$

$$\text{Number of combinations: } C(2+1, 1) = 3!/2!$$

Ex: On a Manhattan street, find # shortest paths from (-1,-2) to (5,8).

$$x_0 + x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = (8 - (-2))$$

Number of combinations:

$$C(10+6, 6) = C(16, 6) = 16!/(10!6!)$$

Integer Linear Equation

Ex: Find the number of integer solutions

$$x+y+z = 20 \text{ with } x \geq 4, y \geq 5, z \geq 6.$$

$$\text{Let } x' = x - 4, y' = y - 5, z' = z - 6,$$

$$\text{i.e. } x = x' + 4, y = y' + 5, z = z' + 6.$$

Thus, we have an equivalent formulation:

$$x' + y' + z' = 5 \text{ with } x', y', z' \geq 0.$$

$$\begin{aligned} \# \text{Combinations: } C(5+2, 2) &= C(7, 2) = \\ 7! / (5!2!) &= (7 \cdot 6) / 2 = 21 \end{aligned}$$

$$\text{Ex: } x+y+z = 20, x < 7, y < 8, z < 9$$

U: no constraints. $|U| = C(20+2, 2)$

A: a set of integer solutions with $x \geq 7$

B: a set of integer solutions with $y \geq 8$

C: a set of integer solutions with $z \geq 9$

A: Let $x' = x - 7$, i.e. $x = x' + 7$ $|A| = C(13+2, 2)$

$$x' + y + z = 13$$

B: Let $y' = y - 8$, i.e. $y = y' + 8$ $|B| = C(12+2, 2)$

$$x + y' + z = 12$$

C: Let $z' = z - 9$ $|C| = C(11+2, 2)$

$$x + y + z' = 11$$

Cont. (from last slide)

$A \cap B: x \geq 7, y \geq 8,$

Let $x' = x - 7, y' = y - 8$

We have $x' + y' + z = 5. \quad |A \cap B| = C(20 - 7 - 8 + 2, 2)$

Likewise, we can derive $|A \cap C| = C(20 - 7 - 9 + 2, 2),$
 $|B \cap C| = C(20 - 8 - 9 + 2, 2), \quad |A \cap B \cap C| = 0.$

According to inclusion and exclusion theorem,
the number of solutions is

$$|U| - |A| - |B| - |C| + |A \cap B| + |A \cap C| + |B \cap C| - |A \cap B \cap C| \\ = 3$$