

# A Pseudopolynomial Algorithm for Alexandrov's Theorem

Daniel Kane, Gregory Nathan Price, and Erik D. Demaine

Massachusetts Institute of Technology

**Abstract.** Alexandrov's Theorem states that every metric with the global topology and local geometry required of a convex polyhedron is in fact the intrinsic metric of some convex polyhedron. Recent work by Bobenko and Izestiev describes a differential equation whose solution is the polyhedron corresponding to a given metric. We describe an algorithm based on this differential equation to compute the polyhedron to arbitrary precision given the metric, and prove a pseudopolynomial bound on its running time.

---

\* G. Price was partially supported by an NSF Graduate Research Fellowship.

\*\* E. Demaine was partially supported by NSF CAREER award CCF-0347776.

# 1 Introduction

Consider the intrinsic metric induced on the surface  $M$  of a convex body in  $\mathbb{R}^3$ . Clearly  $M$  under this metric is homeomorphic to a sphere, and locally convex in the sense that a circle of radius  $r$  has circumference at most  $2\pi r$ .

In 1949, Alexandrov and Pogorelov [1] proved that these two necessary conditions are actually sufficient: every metric space  $M$  that is homeomorphic to a 2-sphere and locally convex can be embedded as the surface of a convex body in  $\mathbb{R}^3$ . Because Alexandrov and Pogorelov's proof is highly nonconstructive, their work opened the question of how to produce the embedding given a concrete  $M$ .

To enable computation we require that  $M$  be a polyhedral metric (space), locally isometric to  $\mathbb{R}^2$  at all but  $n$  points (vertices). Now the theorem is that every polyhedral metric, a complex of triangles with the topology of a sphere and positive curvature at each vertex, can be embedded as an actual convex polyhedron in  $\mathbb{R}^3$ . This case of the Alexandrov-Pogorelov theorem was proven by Alexandrov in 1941 [1], also nonconstructively. Further, Cauchy showed in 1813 [3] that such an embedding must be unique. All the essential geometry of the general case is preserved in the polyhedral case, because every metric satisfying the general hypothesis can be polyhedrally approximated.

In 1996, Sabitov [9, 8, 10, 4] showed how to enumerate all the isometric maps  $M \rightarrow \mathbb{R}^3$  for a polyhedral metric  $M$ , so that one could carry out this enumeration and identify the one map that gives a convex polyhedron. In 2005, Fedorchuk and Pak [5] showed an exponential upper bound on the number of such maps. An exponential lower bound is easy to find, so this algorithm takes time exponential in  $n$  and is therefore unsatisfactory.

Recent work by Bobenko and Izhestiev [2] produced a new proof of Alexandrov's Theorem, describing a certain ordinary differential equation (ODE) and initial conditions whose solution contains sufficient information to construct the embedding by elementary geometry. This work included a computer implementation of the ODE, which empirically produces accurate approximations of embeddings of metrics on which it is tested.

In this work, we describe an algorithm based on the Bobenko-Izhestiev ODE, and prove a pseudopolynomial bound on its running time. Specifically, call an embedding of  $M$   $\varepsilon$ -accurate if the metric is distorted by at most a factor  $1 + \varepsilon$ , and  $\varepsilon$ -convex if each dihedral angle is at most  $\pi + \varepsilon$ . For concreteness,  $M$  may be represented by a list of triangles with side lengths and the names of adjacent triangles. Then we show the following theorem:

**Theorem 1.** *Given a polyhedral metric  $M$  with  $n$  vertices, ratio  $S$  between the diameter and the smallest distance between vertices, and defect (discrete Gaussian curvature) between  $\varepsilon_1$  and  $2\pi - \varepsilon_8$  at each vertex, an  $\varepsilon_6$ -accurate  $\varepsilon_9$ -convex embedding of  $M$  can be found in time  $O(n^{913/2} S^{831} / (\varepsilon_1^{121} \varepsilon_8^{445} \varepsilon_6^{16}))$  where  $\varepsilon = \min(\varepsilon_6/nS, \varepsilon_9\varepsilon_1^2/nS^6)$ .*

The exponents in the time bound of Theorem 1 are remarkably large. Thankfully, no evidence suggests our algorithm actually takes as long to run as the

bound allows. On the contrary, our analysis relies on bounding approximately a dozen geometric quantities, and to keep the analysis tractable we use the simplest bound whenever available. The algorithm’s actual performance is governed by the actual values of these quantities, and therefore by whatever sharper bounds can be proven by a stingier analysis.

To describe our approach, consider an embedding of the metric  $M$  as a convex polyhedron in  $\mathbb{R}^3$ , and choose an arbitrary origin  $O$  in the surface’s interior. Then it is not hard to see that the  $n$  distances  $r_i = \overline{Ov_i}$  from the origin to the vertices  $v_i$ , together with  $M$  and the combinatorial data describing which polygons on  $M$  are faces of the polyhedron, suffice to reconstruct the embedding: the tetrahedron formed by  $O$  and each triangle is rigid in  $\mathbb{R}^3$ , and we have no choice in how to glue them to each other. In Lemma 1 below, we show that in fact the radii alone suffice to reconstruct the embedding, to do so efficiently, and to do so even with radii of finite precision.

Therefore in order to compute the unique embedding of  $M$  that Alexandrov’s Theorem guarantees exists, we compute a set of radii  $r = \{r_i\}_i$  and derive a triangulation  $T$ . The exact radii satisfy three conditions:

1. the radii  $r$  determine nondegenerate tetrahedra from  $O$  to each face of  $T$ ;
2. with these tetrahedra, the dihedral angles at each exterior edge total at most  $\pi$ ; and
3. with these tetrahedra, the dihedral angles about each radius sum to  $2\pi$ .

In our computation, we begin with a set of large initial radii  $r_i = R$  satisfying Conditions 1 and 2, and write  $\kappa = \{\kappa_i\}_i$  for the differences by which Condition 3 fails about each radius. We then iteratively adjust the radii to bring  $\kappa$  near zero and satisfy Condition 3 approximately, maintaining Conditions 1 and 2 throughout.

The computation takes the following form. We describe the Jacobian  $\left(\frac{\partial \kappa_i}{\partial r_j}\right)_{ij}$ , showing that it can be efficiently computed and that its inverse is pseudopolynomially bounded. We show further that the Hessian  $\left(\frac{\partial \kappa_i}{\partial r_j \partial r_k}\right)_{ijk}$  is also pseudopolynomially bounded. It follows that a change in  $r$  in the direction of smaller  $\kappa$  as described by the Jacobian, with some step size only pseudopolynomially small, makes progress in reducing  $|\kappa|$ . The step size can be chosen online by doubling and halving, so it follows that we can take steps of the appropriate size, pseudopolynomial in number, and obtain an  $r$  that zeroes  $\kappa$  to the desired precision in pseudopolynomial total time. Theorem 1 follows.

The construction of [2] is an ODE in the same  $n$  variables  $r_i$ , with a similar starting point and with the derivative of  $r$  driven similarly by a desired path for  $\kappa$ . Their proof differs in that it need only show existence, not a bound, for the Jacobian’s inverse, in order to invoke the inverse function theorem. Similarly, while we must show a pseudopolynomial lower bound (Lemma 14) on the altitudes of the tetrahedra during our computation, the prior work shows only that these altitudes remain positive. In general our computation requires that the known open conditions—this quantity is positive, that map is nondegenerate—be replaced by stronger compact conditions—this quantity is lower-bounded,

that map’s inverse is bounded. We model our proofs of these strengthenings on the proofs in [2] of the simpler open conditions, and we directly employ several other results from that paper where possible.

The remainder of this paper supplies the details of the proof of Theorem 1. We give background in Section 2, and detail the main argument in Section 3. We bound the Jacobian in Section 4 and the Hessian in Section 5. Finally, some lemmas are deferred to Section 6 for clarity.

## 2 Background and Notation

In this section we define our major geometric objects and give the basic facts about them. We also define some parameters describing our central object that we will need to keep bounded throughout the computation.

### 2.1 Geometric notions

Central to our argument are two dual classes of geometric structures introduced by Bobenko and Izestiev in [2] under the names of “generalized convex polytope” and “generalized convex polyhedron”. Because in other usages the distinction between “polyhedron” and “polytope” is that a polyhedron is a three-dimensional polytope, and because both of these objects are three-dimensional, we will refer to these objects as “generalized convex polyhedra” and “generalized convex dual polyhedra” respectively to avoid confusion.

First, we define the objects that our main theorem is about.

**Definition 1.** *A metric  $M$  homeomorphic to the sphere is a polyhedral metric if each  $x \in M$  has an open neighborhood isometric either to a subset of  $\mathbb{R}^2$  or to a cone of angle less than  $2\pi$  with  $x$  mapped to the apex, and if only finitely many  $x$ , called the vertices  $V(M) = \{v_i\}_i$  of  $M$ , fall into the latter case.*

*The defect  $\delta_i$  at a vertex  $v_i \in V(M)$  is the difference between  $2\pi$  and the total angle at the vertex, which is positive by the definition of a vertex.*

*An embedding of  $M$  is a piecewise linear map  $f : M \rightarrow \mathbb{R}^3$ . An embedding  $f$  is  $\varepsilon$ -accurate if it distorts the metric  $M$  by at most  $1 + \varepsilon$ , and  $\varepsilon$ -convex if  $f(M)$  is a polyhedron and each dihedral angle in  $f(M)$  is at most  $\pi + \varepsilon$ .*

*A perfect embedding of a polyhedral metric  $M$  is an isometry  $f : M \rightarrow \mathbb{R}^3$  such that  $f(M)$  is a convex polyhedron. Equivalently, an embedding is perfect if  $0$ -accurate and  $0$ -convex.*

Alexandrov’s Theorem is that every polyhedral metric has a unique perfect embedding, and our contribution is a pseudopolynomial-time algorithm to construct  $\varepsilon$ -accurate  $\varepsilon$ -convex embeddings as approximations to this perfect embedding.

**Definition 2.** *In a tetrahedron  $ABCD$ , write  $\angle CABD$  for the dihedral angle along edge  $AB$ .*

**Definition 3.** A triangulation of a polyhedral metric  $M$  is a decomposition into Euclidean triangles whose vertex set is  $V(M)$ . Its vertices are denoted by  $V(T) = V(M)$ , its edges by  $E(T)$ , and its faces by  $F(T)$ .

A radius assignment on a polyhedral metric  $M$  is a map  $r : V(M) \rightarrow \mathbb{R}_+$ . For brevity we write  $r_i$  for  $r(v_i)$ .

Given a polyhedral metric  $M$ , a triangulation  $T$ , and a radius assignment  $r$ , the generalized convex polyhedron  $P = (M, T, r)$  is a certain metric space on the topological cone on  $M$  with an apex  $O$ , if  $M, T, r$  are suitable. Let the cone  $OF$  for each face  $F \in F(T)$  be isometric to a Euclidean tetrahedron with base  $F$  and side edges given by  $r$ . We require that the total dihedral angle about each edge of  $T$  be at most  $\pi$ , and about each edge  $Ov_i$  at most  $2\pi$ .

Write  $\kappa_i \triangleq 2\pi - \sum_{jk} \angle v_j Ov_i v_k$  for the curvature about  $Ov_i$ , and  $\phi_{ij} \triangleq \angle v_i Ov_j$  for the angle between vertices  $v_i, v_j$  seen from the apex.

Our algorithm, following the construction in [2], will choose a radius assignment for the  $M$  in question and iteratively adjust it until the associated generalized convex polyhedron  $P$  fits nearly isometrically in  $\mathbb{R}^3$ . The resulting radii will give an  $\varepsilon$ -accurate  $\varepsilon$ -convex embedding of  $M$  into  $\mathbb{R}^3$ .

In the argument we will require several geometric objects related to generalized convex polyhedra.

**Definition 4.** A Euclidean simplicial complex is a metric space on a simplicial complex where the metric restricted to each cell is Euclidean.

A generalized convex polygon is a Euclidean simplicial 2-complex homeomorphic to a disk, where all triangles have a common vertex  $V$ , the total angle at  $V$  is no more than  $2\pi$ , and the total angle at each other vertex is no more than  $\pi$ .

Given a generalized convex polyhedron  $P = (M, T, r)$ , the corresponding generalized convex dual polyhedron  $D(P)$  is a certain Euclidean simplicial 3-complex. Let  $O$  be a vertex called the apex,  $A_i$  a vertex with  $OA_i = h_i \triangleq 1/r_i$  for each  $i$ .

For each edge  $v_i v_j \in E(T)$  bounding triangles  $v_i v_j v_k$  and  $v_j v_i v_l$ , construct two simplices  $OA_i A_{jil} A_{ijk}$ ,  $OA_j A_{ijk} A_{jil}$  in  $D(P)$  as follows. Embed the two tetrahedra  $Ov_i v_j v_k, Ov_j v_i v_l$  in  $\mathbb{R}^3$ . For each  $i' \in \{i, j, k, l\}$ , place  $A_{i'}$  along ray  $Ov_{i'}$  at distance  $h_{i'}$ , and draw a perpendicular plane  $P_{i'}$  through the ray at  $A_{i'}$ . Let  $A_{ijk}, A_{jil}$  be the intersection of the planes  $P_i, P_j, P_k$  and  $P_j, P_i, P_l$  respectively.

Now identify the vertices  $A_{ijk}, A_{jki}, A_{kij}$  for each triangle  $v_i v_j v_k \in F(T)$  to produce the Euclidean simplicial 3-complex  $D(P)$ . Since the six simplices produced about each of these vertices  $A_{ijk}$  are all defined by the same three planes  $P_i, P_j, P_k$  with the same relative configuration in  $\mathbb{R}^3$ , the total dihedral angle about each  $OA_{ijk}$  is  $2\pi$ . On the other hand, the total dihedral angle about  $OA_i$  is  $2\pi - \kappa_i$ , and the face about  $A_i$  is a generalized convex polygon of defect  $\kappa_i$ . Let

$$h_{ij} = \frac{h_j - h_i \cos \phi_{ij}}{\sin \phi_{ij}}$$

be the altitude in this face from its apex  $A_i$  to side  $A_{ijk} A_{jil}$ .

**Definition 5.** A spherical simplicial 2-complex is a metric space on a simplicial complex where each 2-cell is isometric to a spherical triangle.

A singular spherical polygon (or triangle, quadrilateral, etc) is a spherical simplicial 2-complex homeomorphic to a disk, where the total angle at each interior vertex is at most  $2\pi$ . A singular spherical polygon is convex if the total angle at each boundary vertex is at most  $\pi$ .

A singular spherical metric is a spherical simplicial 2-complex homeomorphic to a sphere, where the total angle at each vertex is at most  $2\pi$ .

The Jacobian bound in Section 4 makes use of certain multilinear forms described in [2].

**Definition 6.** The dual volume  $\text{vol}(h)$  is the volume of the generalized convex dual polyhedron  $D(P)$ , a cubic form in the dual altitudes  $h$ .

The mixed volume  $\text{vol}(\cdot, \cdot, \cdot)$  is the symmetric trilinear form that formally extends the cubic form  $\text{vol}(\cdot)$ :

$$\text{vol}(a, b, c) \triangleq \frac{1}{6}(\text{vol}(a+b+c) - \text{vol}(a+b) - \text{vol}(b+c) - \text{vol}(c+a) + \text{vol}(a) + \text{vol}(b) + \text{vol}(c)).$$

The  $i$ th dual face area  $E_i(g(i))$  is the area of the face around  $A_i$  in  $D(P)$ , a quadratic form in the altitudes  $g(i) \triangleq \{h_{ij}\}_j$  within this face.

The  $i$ th mixed area  $E_i(\cdot, \cdot)$  is the symmetric bilinear form that formally extends the quadratic form  $E_i(\cdot)$ :

$$E_i(a, b) \triangleq \frac{1}{2}(E_i(a+b) - E_i(a) - E_i(b)).$$

Let  $\pi_i$  be the linear map

$$\pi_i(h)_j \triangleq \frac{h_j - h_i \cos \phi_{ij}}{\sin \phi_{ij}}$$

so that  $\pi_i(h) = g(i)$ . Then define

$$F_i(a, b) \triangleq E_i(\pi_i(a), \pi_i(b)).$$

so that  $F_i(h, h) = E_i(g(i), g(i))$  is the area of face  $i$ .

Observe that  $\text{vol}(h, h, h) = \frac{1}{3} \sum_i h_i F_i(h, h)$ , so that by a simple computation

$$\text{vol}(a, b, c) = \frac{1}{3} \sum_i a_i F_i(b, c).$$

## 2.2 Weighted Delaunay triangulations

The triangulations we require at each step of the computation are the weighted Delaunay triangulations used in the construction of [2]. We give a simpler definition inspired by Definition 14 of [6].

**Definition 7.** The power  $\pi_v(p)$  of a point  $p$  against a vertex  $v$  in a polyhedral metric  $M$  with a radius assignment  $r$  is  $pv^2 - r(v)^2$ .

The center  $C(v_i v_j v_k)$  of a triangle  $v_i v_j v_k \in T(M)$  when embedded in  $\mathbb{R}^2$  is the unique point  $p$  such that  $\pi_{v_i}(p) = \pi_{v_j}(p) = \pi_{v_k}(p)$ , which exists by the radical axis theorem from classical geometry. The quantity  $\pi_{v_i}(p) = \pi(v_i v_j v_k)$  is the power of the triangle.

A triangulation  $T$  of a polyhedral metric  $M$  with radius assignment  $r$  is locally convex at edge  $v_i v_j$  with neighboring triangles  $v_i v_j v_k, v_j v_i v_l$  if  $\pi_{v_l}(C(v_i v_j v_k)) > \pi_{v_l}(v_k)$  and  $\pi_{v_k}(C(v_j v_i v_l)) > \pi_{v_k}(v_l)$  when  $v_i v_j v_k, v_j v_i v_l$  are embedded together in  $\mathbb{R}^2$ .

A weighted Delaunay triangulation for a radius assignment  $r$  on a polyhedral metric  $M$  is a triangulation  $T$  that is locally convex at every edge.

A weighted Delaunay triangulation can be computed in time  $O(n^2 \log n)$  by a simple modification of the “continuous Dijkstra” algorithm of [7].

The radius assignment  $r$  and triangulation  $T$  admits a tetrahedron  $Ov_i v_j v_k$  just if the power of  $v_i v_j v_k$  is negative, and the squared altitude of  $O$  in this tetrahedron is  $-\pi(v_i v_j v_k)$ . The edge  $v_i v_j$  is convex when the two neighboring tetrahedra are embedded in  $\mathbb{R}^3$  just if it is locally convex in the triangulation as in Definition 7. A weighted Delaunay triangulation with negative powers therefore gives a valid generalized convex polyhedron if the curvatures  $\kappa_i$  are positive. For each new radius assignment  $r$  in the computation of Section 3 we therefore compute the weighted Delaunay triangulation and proceed with the resulting generalized convex polyhedron, in which Lemma 14 guarantees a positive altitude and the choices in the computation guarantee positive curvatures.

### 2.3 Notation for bounds

**Definition 8.** Let the following bounds be observed:

1.  $n$  is the number of vertices on  $M$ . By Euler’s formula,  $|E(T)|$  and  $|F(T)|$  are both  $O(n)$ .
2.  $\varepsilon_1 \triangleq \min_i \delta_i$  is the minimum defect.
3.  $\varepsilon_2 \triangleq \min_i (\delta_i - \kappa_i)$  is the minimum defect-curvature gap.
4.  $\varepsilon_3 \triangleq \min_{ij \in E(T)} \phi_{ij}$  is the minimum angle between radii.
5.  $\varepsilon_4 \triangleq \max_i \kappa_i$  is the maximum curvature.
6.  $\varepsilon_5 \triangleq \min_{v_i v_j v_k \in F(T)} \angle v_i v_j v_k$  is the smallest angle in the triangulation. Observe that obtuse angles are also bounded:  $\angle v_i v_j v_k < \pi - \angle v_j v_i v_k \leq \pi - \varepsilon_5$ .
7.  $\varepsilon_6$  is used for the desired accuracy in embedding  $M$ .
8.  $\varepsilon_7 \triangleq (\max_i \frac{\kappa_i}{\delta_i}) / (\min_i \frac{\kappa_i}{\delta_i}) - 1$  is the extent to which the ratio among the  $\kappa_i$  varies from that among the  $\delta_i$ . We will keep  $\varepsilon_7 < \varepsilon_8 / 4\pi$  throughout the computation.
9.  $\varepsilon_8 \triangleq \min_i (2\pi - \delta_i)$  is the minimum angle around a vertex, the complement of the maximum defect.
10.  $\varepsilon_9$  is used for the desired approximation to convexity in embedding  $M$ .

11.  $D$  is the diameter of  $M$ .
12.  $\ell$  is the shortest distance  $v_i v_j$  between vertices.
13.  $S \triangleq D/\ell$  is the maximum ratio of distances.
14.  $d_0 \triangleq \min_{p \in M} Op$  is the minimum height of the apex off of any point on  $M$ .
15.  $d_1 \triangleq \min_{v_i v_j \in E(T)} d(O, v_i v_j)$  is the minimum distance from the apex to any edge of  $T$ .
16.  $d_2 \triangleq \min_i r_i$  is the minimum distance from the apex to any vertex of  $M$ .
17.  $H \triangleq 1/d_0$ ; the name is justified by  $h_i = 1/r_i \leq 1/d_0$ .
18.  $R \triangleq \max_i r_i$ , so  $1/H \leq r_i \leq R$  for all  $i$ .
19.  $T \triangleq HR$  is the maximum ratio of radii.

Of these bounds,  $n, \varepsilon_1, \varepsilon_8, S$  are fundamental to the given metric  $M$ , and  $D$  a dimensionful parameter given by  $M$ . The values  $\varepsilon_6, \varepsilon_9$  define the objective to be achieved, and our computation will drive  $\varepsilon_4$  toward zero while maintaining  $\varepsilon_2$  large and  $\varepsilon_7$  small. In Section 6 we bound the remaining parameters  $\varepsilon_3, \varepsilon_5, R, d_0, d_1, d_2$  in terms of these.

**Definition 9.** Let  $\mathbf{J}$  denote the Jacobian  $(\frac{\partial \kappa_i}{\partial r_j})_{ij}$ , and  $\mathbf{H}$  the Hessian  $(\frac{\partial \kappa_i}{\partial r_j \partial r_k})_{ijk}$ .

### 3 Main Theorem

In this section, we prove our main theorem using the results proved in the remaining sections. Recall

**Theorem 1.** Given a polyhedral metric  $M$  with  $n$  vertices, ratio  $S$  (the spread) between the diameter and the smallest distance between vertices, and defect at least  $\varepsilon_1$  and at most  $2\pi - \varepsilon_8$  at each vertex, an  $\varepsilon_6$ -accurate  $\varepsilon_9$ -convex embedding of  $M$  can be found in time  $O(n^{913/2} S^{831} / (\varepsilon^{121} \varepsilon_1^{445} \varepsilon_8^{616}))$  where  $\varepsilon = \min(\varepsilon_6/nS, \varepsilon_9 \varepsilon_1^2/nS^6)$ .

The algorithm of Theorem 1 obtains an approximate embedding of the polyhedral metric  $M$  in  $\mathbb{R}^3$ . Its main subroutine is described by the following theorem:

**Theorem 2.** Given a polyhedral metric  $M$  with  $n$  vertices, ratio  $S$  (the spread) between the diameter and the smallest distance between vertices, and defect at least  $\varepsilon_1$  and at most  $2\pi - \varepsilon_8$  at each vertex, a radius assignment  $r$  for  $M$  with maximum curvature at most  $\varepsilon$  can be found in time  $O(n^{913/2} S^{831} / (\varepsilon^{121} \varepsilon_1^{445} \varepsilon_8^{616}))$ .

*Proof.* Let a good assignment be a radius assignment  $r$  that satisfies two bounds:  $\varepsilon_7 < \varepsilon_8/4\pi$  so that Lemmas 12–14 apply and  $r$  therefore by the discussion in Subsection 2.2 produces a valid generalized convex polyhedron for  $M$ , and  $\varepsilon_2 = \Omega(\varepsilon_1^2 \varepsilon_8^3/n^2 S^2)$  on which our other bounds rely. By Lemma 9, there exists a good assignment  $r^0$ . We will iteratively adjust  $r^0$  through a sequence  $r^t$  of good

assignments to arrive at an assignment  $r^N$  with maximum curvature  $\varepsilon_4^N < \varepsilon$  as required. At each step we recompute  $T$  as a weighted Delaunay triangulation according to Subsection 2.2.

Given a good assignment  $r = r^n$ , we will compute another good assignment  $r' = r^{n+1}$  with  $\varepsilon_4 - \varepsilon'_4 = \Omega(\varepsilon_1^{445} \varepsilon_4^{121} \varepsilon_8^{616} / (n^{907/2} S^{831}))$ . It follows that from  $r^0$  we can arrive at a satisfactory  $r^N$  with  $N = O((n^{907/2} S^{831}) / (\varepsilon_1^{121} \varepsilon_4^{445} \varepsilon_8^{616}))$ .

To do this, let  $\mathbf{J}$  be the Jacobian  $(\frac{\partial \kappa_i}{\partial r_j})_{ij}$  and  $\mathbf{H}$  the Hessian  $(\frac{\partial^2 \kappa_i}{\partial r_j \partial r_k})_{ijk}$ , evaluated at  $r$ . The goodness conditions and the objective are all in terms of  $\kappa$ , so we choose a desired new curvature vector  $\kappa^*$  in  $\kappa$ -space and apply the inverse Jacobian to get a new radius assignment  $r' = r + \mathbf{J}^{-1}(\kappa^* - \kappa)$  in  $r$ -space. The actual new curvature vector  $\kappa'$  differs from  $\kappa^*$  by an error at most  $\frac{1}{2} |\mathbf{H}| |r' - r|^2 \leq (\frac{1}{2} |\mathbf{H}| |\mathbf{J}^{-1}|^2) |\kappa^* - \kappa|^2$ , quadratic in the desired change in curvatures with a coefficient

$$C \triangleq \frac{1}{2} |\mathbf{H}| |\mathbf{J}^{-1}|^2 = O\left(\frac{n^{3/2} S^{14}}{\varepsilon_5^3} \frac{R^{23}}{D^{14} d_0^3 d_1^8} \left(\frac{n^{7/2} T^2}{\varepsilon_2 \varepsilon_3 \varepsilon_4} R\right)^2\right) = O\left(\frac{n^{905/2} S^{831}}{\varepsilon_1^{443} \varepsilon_4^{121} \varepsilon_8^{616}}\right)$$

by Theorems 3 and 4 and Lemmas 10, 9, 14, and 11.

Therefore pick a step size  $p$ , and choose  $\kappa^*$  according to  $\kappa_i^* - \kappa_i = -p\kappa_i - p(\kappa_i - \delta_i \min_j \frac{\kappa_j}{\delta_j})$ . The first term diminishes all the curvatures together to reduce  $\varepsilon_4$ , and the second rebalances them to keep the ratios  $\frac{\kappa_j}{\delta_j}$  nearly equal so that  $\varepsilon_7$  remains small. In Subsection A.1 in the Appendix we show that the resulting actual curvatures  $\kappa'$  make  $r'$  a good assignment and put  $\varepsilon'_4 \leq \varepsilon_4 - p\varepsilon_4/2$ , so long as

$$p \leq \varepsilon_1^2 / 64\pi^2 n \varepsilon_4 C. \quad (1)$$

This produces a good radius assignment  $r'$  in which  $\varepsilon_4$  has declined by at least

$$\frac{p\varepsilon_4}{2} = \frac{\varepsilon_1^2}{128\pi^2 n C} = \Omega\left(\frac{\varepsilon_1^{445} \varepsilon_4^{121} \varepsilon_8^{616}}{n^{907/2} S^{831}}\right)$$

as required.

As a simplification, we need not compute  $p$  exactly according to (1). Rather, we choose the step size  $p^t$  at each step, trying first  $p^{t-1}$  (with  $p^0$  an arbitrary constant) and computing the actual curvature error  $|\kappa' - \kappa^*|$ . If the error exceeds its maximum acceptable value  $p\varepsilon_1^2 \varepsilon_4 / 16\pi^2$  then we halve  $p^t$  and try step  $t$  again, and if it falls below half this value then we double  $p^t$  for the next round. Since we double at most once per step and halve at most once per doubling plus a logarithmic number of times to reach an acceptable  $p$ , this doubling and halving costs only a constant factor. Even more important than the resulting simplification of the algorithm, this technique holds out the hope of actual performance exceeding the proven bounds.

Now each of the  $N$  iterations of the computation go as follows. Compute the weighted Delaunay triangulation  $T^t$  for  $r^t$  in time  $O(n^2 \log n)$  as described in Subsection 2.2. Compute the Jacobian  $\mathbf{J}^t$  in time  $O(n^2)$  using formulas (14, 15) in [2]. Choose a step size  $p^t$ , possibly adjusting it, as discussed above. Finally, take the resulting  $r'$  as  $r^{t+1}$  and continue. The computation of  $\kappa^*$  to check  $p^t$  runs in linear time, and that of  $r'$  in time  $O(n^\omega)$  where  $\omega < 3$  is the time exponent of matrix multiplication. Each iteration therefore costs time  $O(n^3)$ , and the whole computation costs time  $O(n^3 N)$  as claimed.

Now with our radius assignment  $r$  for  $M$  and the resulting generalized convex polyhedron  $P$  with curvatures all near zero, it remains to approximately embed  $P$  and therefore  $M$  in  $\mathbb{R}^3$ . To begin, we observe that this is easy to do given exact values for  $r$  and in a model with exact computation: after triangulating,  $P$  is made up of rigid tetrahedra and we embed one tetrahedron arbitrarily, then embed each neighboring tetrahedron in turn.

In a realistic model, we compute only with bounded precision, and in any case Theorem 2 gives us only curvatures near zero, not equal to zero. Lemma 1 produces an embedding in this case, settling for less than exact isometry and exact convexity.

**Lemma 1.** *There is an algorithm that, given a radius assignment  $r$  for which the corresponding curvatures  $\kappa_i$  are all less than  $\varepsilon = O(\min(\varepsilon_6/nS, \varepsilon_9\varepsilon_1^2/nS^6))$  for some constant factor, produces explicitly by vertex coordinates in time  $O(n^2 \log n)$  an  $\varepsilon_6$ -accurate  $\varepsilon_9$ -convex embedding of  $M$ .*

*Proof (sketch).* For details see Subsection A.1.

As in the exact case, triangulate  $M$ , embed one tetrahedron arbitrarily, and then embed its neighbors successively. The positive curvature will force gaps between the tetrahedra. Then replace the several copies of each vertex by their centroid, so that the tetrahedra are distorted but leave no gaps. This is the desired embedding. The proofs of  $\varepsilon_6$ -accuracy and  $\varepsilon_9$ -convexity are straightforward and deferred to Subsection A.1.

A weighted Delaunay triangulation takes time  $O(n^2 \log n)$  as discussed in Subsection 2.2, and the remaining steps take time  $O(n)$ .

We now have all the pieces to prove our main theorem.

*Proof (Theorem 1).* Let  $\varepsilon \triangleq O(\min(\varepsilon_6/nS, \varepsilon_9\varepsilon_1^2/nS^6))$ , and apply the algorithm of Theorem 2 to obtain in time  $O(n^{913/2} S^{831} / (\varepsilon^{121} \varepsilon_1^{445} \varepsilon_8^{616}))$  a radius assignment  $r$  for  $M$  with maximum curvature  $\varepsilon_4 \leq \varepsilon$ .

Now apply the algorithm of Lemma 1 to obtain in time  $O(n^2 \log n)$  the desired embedding and complete the computation.

## 4 Bounding the Jacobian

**Theorem 3.** *The Jacobian  $\mathbf{J} = \left(\frac{\partial \kappa_i}{\partial r_j}\right)_{ij}$  has inverse pseudopolynomially bounded by  $|\mathbf{J}^{-1}| = O\left(\frac{n^{7/2}T^2}{\varepsilon_2\varepsilon_3^3\varepsilon_4}R\right)$ .*

*Proof.* Our argument parallels that of Corollary 2 in [2], which concludes that the same Jacobian is nondegenerate. Theorem 4 of [2] shows that this Jacobian equals the Hessian of the volume of the dual  $D(P)$ . The meat of the corollary's proof is in Theorem 5 of [2], which begins by equating this Hessian to the bilinear form  $6 \text{vol}(h, \cdot, \cdot)$  derived from the mixed volume we defined in Definition 6. So we have to bound the inverse of this bilinear form.

To do this it suffices to show that the form  $\text{vol}(h, x, \cdot)$  has norm at least  $\Omega\left(\frac{\varepsilon_2\varepsilon_3^3\varepsilon_4}{n^{7/2}T^2} \frac{|x|}{R}\right)$  for all vectors  $x$ . Equivalently, suppose some  $x$  has  $|\text{vol}(h, x, z)| \leq |z|$  for all  $z$ ; we show  $|x| = O\left(\frac{n^{7/2}T^2}{\varepsilon_2\varepsilon_3^3\varepsilon_4}R\right)$ .

To do this we follow the proof in Theorem 5 of [2] that the same form  $\text{vol}(h, x, \cdot)$  is nonzero for  $x$  nonzero. Throughout the argument we work in terms of the dual  $D(P)$ .

Recall that for each  $i$ ,  $\pi_i x$  is defined as the vector  $\{x_{ij}\}_j$ . It suffices to show that for all  $i$

$$|\pi_i x|_2^2 = O\left(\frac{n^3 T^3}{\varepsilon_2^2 \varepsilon_3 \varepsilon_4} R^2 + \frac{n^2 T^2}{\varepsilon_2 \varepsilon_3 \varepsilon_4} R |x|_1\right)$$

since then by Lemma 2

$$|x|_2^2 \leq \frac{4n}{\varepsilon_3^2} \max_i |\pi_i x|_2^2 = O\left(\frac{n^4 T^3}{\varepsilon_2^2 \varepsilon_3^3 \varepsilon_4} R^2 + \frac{n^3 T^2}{\varepsilon_2 \varepsilon_3 \varepsilon_4} R |x|_1\right),$$

and since  $|x|_1 \leq \sqrt{n}|x|_2$  and  $X^2 \leq a + bX$  implies  $X \leq \sqrt{a} + b$ ,  $|x|_2 = O\left(\frac{n^{7/2}T^2}{\varepsilon_2\varepsilon_3^3\varepsilon_4}R\right)$ . Therefore fix an arbitrary  $i$ , let  $g = \pi_i h$  and  $y = \pi_i x$ , and we proceed to bound  $|y|_2$ .

We break the space on which  $E_i$  acts into the 1-dimensional positive eigenspace of  $E_i$  and its  $(k-1)$ -dimensional negative eigenspace, since by Lemma 3.4 of [2] the signature of  $E_i$  is  $(1, k-1)$ , where  $k$  is the number of neighbors of  $v_i$ . Write  $\lambda_+$  for the positive eigenvalue and  $-E_i^-$  for the restriction to the negative eigenspace so that  $E_i^-$  is positive definite, and decompose  $g = g_+ + g_-$ ,

$y = y_+ + y_-$  by projection into these subspaces. Then we have

$$\begin{aligned} G &\triangleq E_i(g, g) = \lambda_+ g_+^2 - E_i^-(g_-, g_-) \triangleq \lambda_+ g_+^2 - G_- \\ E_i(g, y) &= \lambda_+ g_+ y_+ - E_i^-(g_-, y_-) \\ Y &\triangleq E_i(y, y) = \lambda_+ y_+^2 - E_i^-(y_-, y_-) \triangleq \lambda_+ y_+^2 - Y_- \end{aligned}$$

and our task is to obtain an upper bound on  $Y_- = E_i^-(y_-, y_-)$ , which will translate through our bound on the eigenvalues of  $E_i$  away from zero into the desired bound on  $|y|$ .

We begin by obtaining bounds on  $|E_i(g, y)|$ ,  $G_-$ ,  $G$ , and  $Y$ . Since  $|z| \geq |\text{vol}(h, x, z)|$  for all  $z$  and  $\text{vol}(h, x, z) = \sum_j z_j F_j(h, x)$ , we have  $|E_i(g, y)| = |F_i(h, x)| \leq 1$ . Further,  $\det \begin{pmatrix} E_i(g, g) & E_i(y, g) \\ E_i(g, y) & E_i(y, y) \end{pmatrix} < 0$  because  $E_i$  has signature  $(1, 1)$  restricted to the  $(y, g)$  plane, so by Lemma 3  $Y = E_i(y, y) < \frac{R^2}{\varepsilon_2}$ . On the other hand  $-|x|_1 < \sum_j x_j F_j(x, h) = \sum_j h_j F_j(x, x)$ , so

$$Y = E_i(y, y) = F_i(x, x) > -\frac{1}{h_i} \left( (n-1) \frac{R^2}{\varepsilon_2} H + |x|_1 \right) > -\left( \frac{nT}{\varepsilon_2} R^2 + R|x|_1 \right).$$

Now  $G = E_i(g, g) > 0$ , being the area of the face about  $A_i$  in  $D(P)$ . We have  $|E_i| = O(n/\varepsilon_3)$  by construction, so  $G, G_- \leq G + G_- \leq |E_i| |h|^2 = O(nH^2/\varepsilon_3)$  and similarly  $G = O(nH^2/\varepsilon_3)$ . On the other hand we have  $G = \Omega(\varepsilon_2/R^2)$  by Lemma 3.

Now, observe that  $\lambda_+ y_+ g_+$  is the geometric mean  $\sqrt{(\lambda_+ g_+^2)(\lambda_+ y_+^2)} = \sqrt{(G + G_-)(Y + Y_-)}$  and by Cauchy-Schwarz  $E_i^-(y_-, g_-) \leq \sqrt{G_- Y_-}$ , so that

$$\begin{aligned} 1 &\geq E_i(y, g) \geq \sqrt{(G + G_-)(Y + Y_-)} - \sqrt{G_- Y_-} \\ &= \sqrt{Y_-} \frac{G}{\sqrt{G + G_-} + \sqrt{G_-}} + \sqrt{G + G_-} \frac{Y}{\sqrt{Y + Y_-} + \sqrt{Y_-}}. \end{aligned}$$

By elementary algebra, analyzing separately the cases  $Y \geq 0, Y < 0$ , it follows that

$$Y_- = O\left( \frac{n^2 T^3}{\varepsilon_2^2 \varepsilon_3} R^2 + \frac{n T^2}{\varepsilon_2 \varepsilon_3} R|x|_1 \right).$$

Then using  $Y \leq R/\varepsilon_2^2$  and Lemma 4 we have

$$|y|_2^2 = y_+^2 + |y_-|_2^2 \leq |E_i^{-1}|((Y + Y_-) + Y_-) = O\left( \frac{n^3 T^3}{\varepsilon_2^2 \varepsilon_3 \varepsilon_4} R^2 + \frac{n^2 T^2}{\varepsilon_2 \varepsilon_3 \varepsilon_4} R|x|_1 \right)$$

and the theorem follows.

Three small lemmas used above follow from the geometry of singular spherical polygons and of generalized convex dual polyhedra. Their proofs are deferred to Subsection A.2 in the Appendix for brevity.

**Lemma 2.**  $|x|^2 \leq (4n/\varepsilon_3^2) \max_i |\pi_i x|^2$ .

**Lemma 3.**  $F_i(h, h) > \varepsilon_2/R^2$ .

**Lemma 4.** *The inverse of the form  $E_i$  is bounded by  $|E_i^{-1}| = O(n/\varepsilon_4)$ .*

## 5 Bounding the Hessian

In order to control the error in each step of our computation, we need to keep the Jacobian  $\mathbf{J}$  along the whole step close to the value it started at, on which the step was based. To do this we bound the Hessian  $\mathbf{H}$  when the triangulation is fixed, and we show that the Jacobian does not change discontinuously when changing radii force a new triangulation.

Each curvature  $\kappa_i$  is of the form  $2\pi - \sum_{j,k:v_i v_j v_k \in T} \angle v_j O v_i v_k$ , so in analyzing its derivatives we focus on the dihedral angles  $\angle v_j O v_i v_k$ . When the tetrahedron  $O v_i v_j v_k$  is embedded in  $\mathbb{R}^3$ , the angle  $\angle v_j O v_i v_k$  is determined by elementary geometry as a smooth function of the distances among  $O, v_i, v_j, v_k$ . For a given triangulation  $T$  this makes  $\kappa$  a smooth function of  $r$ . Our first lemma shows that no error is introduced at the transitions where the triangulation  $T(r)$  changes.

**Lemma 5.** *The Jacobian  $\mathbf{J} = \left(\frac{\partial \kappa_i}{\partial r_j}\right)_{ij}$  is continuous at the boundary between radii corresponding to one triangulation and to another.*

*Proof (sketch).* For a full proof, see Subsection A.3. The proof uses elementary geometry to compare the figures determined by two triangulations near a radius assignment on their boundary.

It now remains to control the change in  $\mathbf{J}$  as  $r$  changes within any particular triangulation, which we do by bounding the Hessian.

**Theorem 4.** *The Hessian  $\mathbf{H} = \left(\frac{\partial \kappa_i}{\partial r_j \partial r_k}\right)_{ijk}$  has norm  $O(n^{5/2} S^{14} R^{23} / (\varepsilon_5^3 d_0^3 d_1^8 D^{14}))$ .*

*Proof.* It suffices to bound in absolute value each element  $\frac{\partial^2 \kappa_i}{\partial r_j \partial r_k}$  of the Hessian. Since  $\kappa_i$  is  $2\pi$  minus the sum of the dihedral angles about radius  $r_i$ , its derivatives decompose into sums of derivatives  $\frac{\partial^2 \angle v_l O v_i v_m}{\partial r_j \partial r_k}$  where  $v_i v_l v_m \in F(T)$ . Since the geometry of each tetrahedron  $O v_i v_l v_m$  is determined by its own side lengths, the only nonzero terms are where  $j, k \in \{i, l, m\}$ .

It therefore suffices to bound the second partial derivatives of dihedral angle  $AB$  in a tetrahedron  $ABCD$  with respect to the lengths  $AB, AC, AD$ . By Lemma 8 below, these are degree-23 polynomials in the side lengths of  $ABCD$ , divided by  $[ABCD]^3[ABC]^4[ABD]^4$ . Since  $2[ABC], 2[ABD] \geq (D/S)d_1$ ,  $6[ABCD] \geq d_0(D/S)^2 \sin \varepsilon_5$ , and each side is  $O(R)$ , the second derivative is  $O(S^{14}R^{23}/(\varepsilon_5^3 d_0^3 d_1^8 D^{14}))$ .

Now each element in the Hessian is the sum of at most  $n$  of these one-tetrahedron derivatives  $\frac{\partial^2 \angle v_i O v_i v_m}{\partial r_j \partial r_k}$ , and the norm of the Hessian itself is at most  $n^{3/2}$  times the greatest absolute value of any of its elements, so the theorem is proved.

**Definition 10.** *For the remainder of this section,  $ABCD$  is a tetrahedron and  $\theta$  the dihedral angle  $\angle CABD$  on  $AB$ .*

**Lemma 6.**  $\sin \theta = \frac{3}{2} \frac{[ABCD][AB]}{[ABC][ABD]}$ .

*Proof (sketch).* Elementary geometry; see Subsection A.3 for details.

**Lemma 7.** *Each of the derivatives  $\frac{\partial \theta}{\partial AB}, \frac{\partial \theta}{\partial AC}, \frac{\partial \theta}{\partial AD}$  is a degree-10 polynomial in the side lengths of  $ABCD$ , divided by  $[ABCD][ABC]^2[ABD]^2$ .*

*Proof (sketch).* See Subsection A.3 for a full proof. We write  $[ABC]^2, [ABD]^2, [ABCD]^2$  as polynomials in the side lengths and apply Lemma 6 to obtain  $\sin^2 \theta$  as a rational function of the side lengths. Then elementary calculus, followed by some grungy but straightforward algebra that may be done in SAGE [11] or another computer algebra system, proves the lemma.

**Lemma 8.** *Each of the six second partial derivatives of  $\theta$  in  $AB, AC, AD$  is a degree-23 polynomial in the side lengths of  $ABCD$ , divided by  $[ABCD]^3[ABC]^4[ABD]^4$ .*

*Proof (sketch).* Lemma 7 followed by straightforward calculus; see Subsection A.3 for details.

## 6 Intermediate Bounds

In this section we bound miscellaneous parameters in the computation in terms of the fundamental parameters  $n, S, \varepsilon_1, \varepsilon_8$  and the computation-driving parameter  $\varepsilon_4$ .

## 6.1 Initial conditions

**Lemma 9.** *Given a polyhedral metric space  $M$ , there exists a radius assignment  $r$  with curvature skew  $\varepsilon_7 < \varepsilon_8/4\pi$ , maximum radius  $R = O(nD/\varepsilon_1\varepsilon_8)$ , and minimum defect-curvature gap  $\varepsilon_2 = \Omega(\varepsilon_1^2\varepsilon_8^3/n^2S^2)$ .*

*Proof (sketch).* Take  $r_i = R$  for all  $i$ , with  $R$  sufficiently large. Then each  $\kappa_i$  is nearly equal to  $\delta_i$ , so that  $\varepsilon_7$  is small. For the quantitative bounds and a full proof, see Subsection A.4 in the Appendix.

## 6.2 Two angle bounds

These bounds can be proven by elementary geometry; details deferred to Subsection A.5 in the Appendix for brevity.

**Lemma 10.**  $\varepsilon_3 > \ell d_1/R^2$ .

**Lemma 11.**  $\varepsilon_5 > \varepsilon_2/6S$ .

## 6.3 Keeping away from the surface

In this section we bound  $O$  away from the surface  $M$ . The bounds are effective versions of Lemmas 4.8, 4.6, and 4.5 respectively of [2], and the proofs, deferred for brevity to Subsection A.6 in the Appendix, are similar but more involved.

Recall that  $d_2$  is the minimum distance from  $O$  to any vertex of  $M$ ,  $d_1$  is the minimum distance to any edge of  $T$ , and  $d$  is the minimum distance from  $O$  to any point of  $M$ .

**Lemma 12.**  $d_2 = \Omega(D\varepsilon_1\varepsilon_4\varepsilon_5^2\varepsilon_8/(nS^4))$ .

**Lemma 13.**  $d_1 = \Omega(D\varepsilon_1^2\varepsilon_4^4\varepsilon_5^6\varepsilon_8^2/(n^2S^{10}))$ .

**Lemma 14.**  $d_0 = \Omega(D\varepsilon_1^4\varepsilon_4^9\varepsilon_5^{12}\varepsilon_8^4/(n^4S^{22}))$ .

## References

1. A. D. Alexandrov. *Convex Polyhedra*. Springer-Verlag, Berlin, 2005.
2. A. I. Bobenko and I. Izmistiev. Alexandrov's theorem, weighted Delaunay triangulations, and mixed volumes. *Annales de l'Institut Fourier*, 2006, arXiv:math.DG/0609447.
3. A. L. Cauchy. Sur les polygones et les polyèdres, seconde mémoire. *J. École Polytechnique*, XVIe Cahier, Tome IX:113–148, 1813. In *OEuvres Complètes*, IIe Série, volume 1, pages 26–38, Paris, 1905.
4. E. D. Demaine and J. O'Rourke. *Geometric Folding Algorithms*. Cambridge University Press, Cambridge, 2007.
5. M. Fedorchuk and I. Pak. Rigidity and polynomial invariants of convex polytopes. *Duke Math. J.*, 129:371–404, 2005.
6. David Glickenstein. Geometric triangulations and discrete Laplacians on manifolds, 2005, arXiv:math/0508188v1.
7. J. S. B. Mitchell, D. M. Mount, and C. H. Papadimitriou. The discrete geodesic problem. *SIAM J. Comput.*, 16:647–668, 1987.
8. I. Sabitov. The volume of a polyhedron as a function of its metric. *Fundam. Prikl. Mat.*, 2(4):1235–1246, 1996.
9. I. Sabitov. The volume of a polyhedron as a function of its metric and algorithmical solution of the main problems in the metric theory of polyhedra. In *International School-Seminar Devoted to the N. V. Efimov's Memory*, pages 64–65, 1996.
10. I. Sabitov. The volume as a metric invariant of polyhedra. *Discrete Comput. Geom.*, 20:405–425, 1998.
11. W. Stein. *Sage: Open Source Mathematical Software (version 2.10.1)*. The Sage Group, 2008. <http://www.sagemath.org>.

## A Additional proofs

### A.1 Proofs for Section 3

In this section we give additional details to complete the proof of Theorem 2, and a complete proof of Lemma 1.

*Proof (Proof for part of Theorem 2).* In the proof of Theorem 2, after we pick a step size  $p$  we choose  $\kappa^*$  according to

$$\kappa_i^* - \kappa_i = -p\kappa_i - p \left( \kappa_i - \delta_i \min_j \frac{\kappa_j}{\delta_j} \right). \quad (2)$$

Here we prove the assertion in the main text that the resulting actual curvatures  $\kappa'$  make  $r'$  a good assignment and put  $\varepsilon'_4 \leq \varepsilon_4 - p\varepsilon_4/2$ , so long as

$$p \leq \varepsilon_1^2 / 64\pi^2 n \varepsilon_4 C. \quad (3)$$

Consider a hypothetical  $r^*$  that gives the curvatures  $\kappa^*$ , and examine the conditions on  $\varepsilon_4, \varepsilon_2, \varepsilon_7$  in turn.

Both terms on the right-hand side of (2) are nonpositive, so each  $\kappa_i$  decreases by at least  $p\kappa_i$ . Therefore the maximum curvature  $\varepsilon_4$  decreases by at least  $p\varepsilon_4$ . If any defect-curvature gap  $\delta_i - \kappa_i$  is less than  $\varepsilon_1/2$ , then it increases by at least  $p\kappa_i \geq p(\delta_i - \varepsilon_1/2) \geq p(\varepsilon_1/2)$ ; so the minimum defect-curvature gap  $\varepsilon_2$  either increases by at least  $p\varepsilon_1/2$  or is at least  $\varepsilon_1/2$  already. Finally, the  $-p\kappa_i$  term decreases each  $\kappa_i$  in the same ratio and therefore preserves  $\varepsilon_7$ , and the  $-p(\kappa_i - \delta_i \min_j(\kappa_j/\delta_j))$  term decreases each ratio  $\kappa_i/\delta_i$  by  $p$  times the difference  $(\kappa_i/\delta_i - \min_j(\kappa_j/\delta_j))$  and therefore reduces  $\varepsilon_7$  by  $p\varepsilon_7$ . Therefore  $\kappa^*$  would satisfy all three conditions with some room to spare.

In particular, if we choose  $p$  to guarantee that each  $\kappa'_i$  differs from  $\kappa_i^*$  by at most  $p\varepsilon_4/2$ , at most  $p\varepsilon_1/2$ , and at most  $p(\varepsilon_1/4\pi)(\min_i \kappa_i)$ , then this discussion shows that the step from  $r$  to  $r'$  will make at least half the ideal progress  $p\varepsilon_4$  in  $\varepsilon_4$  and keep  $\varepsilon_2, \varepsilon_7$  within bounds.

Now

$$\min_i \kappa_i \geq (\max_j \kappa_j) (\min_{ij} \delta_i/\delta_j) (1 + \varepsilon_7)^{-1} \geq \varepsilon_4 (\varepsilon_1/2\pi)/2 = \varepsilon_1 \varepsilon_4 / 4\pi$$

and

$$|\kappa' - \kappa^*|_\infty \leq |\kappa' - \kappa^*| \leq C |\kappa^* - \kappa|^2 \leq 4C p^2 |\kappa|^2 \leq 4C p^2 n \varepsilon_4^2$$

and it follows that choosing  $p$  according to (3) suffices to ensure that  $r'$  is a good assignment and  $\varepsilon_4$  declines by at least  $p\varepsilon_4/2$ .

**Lemma 1.** *There is an algorithm that, given a radius assignment  $r$  for which the corresponding curvatures  $\kappa_i$  are all less than  $\varepsilon = O(\min(\varepsilon_6/nS, \varepsilon_9\varepsilon_1^2/nS^6))$  for some constant factor, produces explicitly by vertex coordinates in time  $O(n^2 \log n)$  an  $\varepsilon_6$ -accurate  $\varepsilon_9$ -convex embedding of  $M$ .*

*Proof.* Compute the weighted Delaunay triangulation  $T$  of  $r$  on  $M$ , and consider the tetrahedra  $OF_i$  for  $F \in F(T)$ . Embed some  $OF_i$  in  $\mathbb{R}^3$  arbitrarily, embed its neighbors next to it, and so forth, leaving gaps as required by the positive curvature, and call this configuration  $Q$ . Since the curvature around each radius is less than  $\varepsilon$ , the several copies of each vertex will be separated by at most  $n\varepsilon D$ . Now replace the several copies of each vertex by their centroid, so that the tetrahedra are distorted but leave no gaps. Call the resulting polyhedron  $P$  and its surface metric  $M'$ . The computation of the weighted Delaunay triangulation takes time  $O(n^2 \log n)$  as discussed in Subsection 2.2, and the remaining steps require time  $O(n)$ . We claim this embedding is  $\varepsilon_6$ -accurate and  $\varepsilon_9$ -convex.

To show  $\varepsilon_6$ -accuracy, observe that since each copy of each vertex was moved by at most  $n\varepsilon D$  from  $Q$  to  $P$ , no edge of any triangle was stretched by more than a ratio  $n\varepsilon S$ , and the piecewise linear map between faces relates  $M'$  to  $M$  with distortion  $n\varepsilon S \leq \varepsilon_6$  as required.

Now we show  $\varepsilon_9$ -convexity. Consider two neighboring triangles  $v_i v_j v_k, v_j v_i v_l$  in  $T$ ; we will show the exterior dihedral angle is at least  $-\varepsilon_9$ . First, consider repeating the embedding with  $Ov_i v_j v_k$  the original tetrahedron, so that  $Ov_i v_j v_k, Ov_j v_i v_l$  embed without gaps. This moves each vertex by at most  $n\varepsilon D$ , and makes the angle  $v_l v_i v_j v_k$  convex and the tetrahedron  $v_l v_i v_j v_k$  have positive signed volume. The volume of this tetrahedron in the  $P$  configuration is therefore at least  $-n\varepsilon D^3$ , since the derivative of the volume in any vertex is the area of the opposite face, which is at always less than  $D^2$  since the sides remain  $(1 + o(1))D$  in length.

Therefore suppose the exterior angle  $\angle v_l v_i v_j v_k$  is negative. Then by Lemma 6 and Lemma 11,

$$\sin \angle v_l v_i v_j v_k = \frac{3 [v_l v_i v_j v_k][v_i v_j]}{2 [v_i v_j v_l][v_j v_i v_k]} \geq -\frac{(n\varepsilon D^3)D}{(\ell^2 \varepsilon_5/4)^2} \geq -\varepsilon \frac{576nS^6}{\varepsilon_2^2}$$

and since  $\varepsilon_2 \geq \varepsilon_1/2$  at the end of the computation,  $\angle v_l v_i v_j v_k \geq -\varepsilon 2304nS^6/\varepsilon_1^2 \geq -\varepsilon_9$  as claimed.

## A.2 Proofs for Section 4

In this subsection we prove three ancillary lemmas from Section 4.

**Lemma 2.**  $|x|^2 \leq (4n/\varepsilon_3^2) \max_i |\pi_i x|^2$ .

*Proof.* Let  $i = \arg \max_i |x_i|$ , and let  $v_j$  be a neighbor in  $T$  of  $v_i$ . Without loss of generality let  $x_i > 0$ . Then

$$(\pi_j x)_i = \frac{x_i - x_j \cos \phi_{ij}}{\sin \phi_{ij}} \geq x_i \frac{1 - \cos \phi_{ij}}{\sin \phi_{ij}} = x_i \tan(\phi_{ij}/2) > x_i \phi_{ij}/2 \geq |x|_\infty \varepsilon_3/2$$

and it follows that

$$|\pi_i x| \geq |\pi_i x|_\infty > |x|_\infty \varepsilon_3/2 \geq |x|_\infty \varepsilon_3/2\sqrt{n}$$

which proves the lemma.

**Lemma 3.**  $F_i(h, h) > \varepsilon_2/R^2$ .

*Proof.* The proof of Proposition 8 in [2] shows that a certain singular spherical polygon has angular area  $\delta_i - \kappa_i$ , where the singular spherical polygon is obtained by stereographic projection of each simplex of  $P_i^*$  onto a sphere of radius  $1/r_i$  tangent to it. The total area of the polygon is  $(\delta_i - \kappa_i)/r_i^2$  at this radius, so because projection of a plane figure onto a tangent sphere only decreases area we have  $F_i(h, h) = \text{area}(P_i^*) > (\delta_i - \kappa_i)/r_i^2 > \varepsilon_2/R^2$ .

**Lemma 4.** *The inverse of the form  $E_i$  is bounded by  $|E_i^{-1}| = O(n/\varepsilon_4)$ .*

*Proof.* We follow the argument in Lemma 3.4 of [2] that the same form is non-degenerate. Let  $\ell_j(y)$  be the length of the side between  $A_i$  and  $A_j$  in  $D(P)$  when the altitudes  $h_{ij}$  are given by  $y$ . Since  $E_i(y) = \frac{1}{2} \sum_j \ell_j(y) y_j$  it follows that  $E_i(a, b) = \frac{1}{2} \sum_j \ell_j(a) b_j$ . Therefore in order to bound the inverse of the form  $E_i$  it suffices to bound the inverse of the linear map  $\ell$ .

Consider a  $y$  such that  $|\ell(y)|_\infty \leq 1$ ; we will show  $|y|_\infty = O(n/\varepsilon_4)$ . Unfold the generalized polygon described by  $y$  into the plane, apex at the origin; the sides are of length  $\ell_j(y)$ , so the first and last vertex are a distance at most  $|\ell(y)|_1 \leq n$  from each other. But the sum of the angles is at least  $\varepsilon_4$  short of  $2\pi$ , so this means all the vertices are within  $O(n/\varepsilon_4)$  of the origin; and the altitudes  $y_j$  are no more than the distances from vertices to the origin, so they are also  $O(n/\varepsilon_4)$  as claimed.

### A.3 Proofs for Section 5

In this subsection we give details for the proof sketches in Section 5.

**Lemma 5.** *The Jacobian  $\mathbf{J}$  is continuous at the boundary between radii corresponding to one triangulation and to another.*

*Proof.* Let  $r$  be a radius assignment consistent with more than one triangulation, say with a flat face  $v_i v_j v_k v_l$  that can be triangulated by  $v_i v_k$  as  $v_i v_j v_k, v_k v_l v_i$  or by  $v_j v_l$  as  $v_j v_k v_l, v_l v_i v_j$ . Since the Jacobian is continuous when either triangulation is fixed and  $r$  varies, it suffices to show that for neighboring radius assignments  $r + \Delta r$ , the curvatures  $\kappa$  obtained with either triangulation differ by a magnitude  $O(|\Delta r|^2)$ , with any coefficient determined by the polyhedral metric or the radius assignment  $r$ .

Embed the two tetrahedra  $Ov_i v_j v_k, Ov_k v_l v_i$  or  $Ov_j v_k v_l, Ov_l v_i v_j$  together in  $\mathbb{R}^3$ , with distances  $[Ov_i]$ , etc., taken from  $r + \Delta r$ . Of the ten pairwise distances between the five points in this diagram, eight are determined by  $M$  or the radii and do not vary between the  $v_i v_k$  and  $v_j v_l$  diagrams. Since the angles  $\angle v_j O v_i v_k$ , etc., are smooth functions of these ten distances, it suffices to show that the remaining two distances  $[v_i v_k], [v_j v_l]$  differ between the diagrams by  $O(|\Delta r|^2)$ . Letting  $X$  denote the intersection of the geodesics  $v_i v_k, v_j v_l$  on the face  $v_i v_j v_k v_l$ , we have  $[v_i v_k]$  in the  $v_i v_k$  diagram equal to  $[v_i X] + [X v_k]$ , while in the  $v_j v_l$  diagram  $v_i X v_k$  form a triangle with the same lengths  $[v_i X], [X v_k]$  and a shorter  $[v_k v_i]$ . The difference between  $[v_i v_k]$  in the two diagrams is therefore the slack in the triangle inequality in this triangle  $v_i X v_k$ , which is bounded by  $O(|\Delta r|^2)$  since the vertices have moved a distance  $O(|\Delta r|)$  from where  $r$  placed them with  $v_i, X, v_k$  collinear.

**Lemma 6.**

$$\sin \theta = \frac{3 [ABCD][AB]}{2 [ABC][ABD]}.$$

*Proof.* First, translate  $C$  and  $D$  parallel to  $AB$  to make  $BCD$  perpendicular to  $AB$ , which has no effect on either side of the equation. Now  $[ABCD] = [BCD][AB]/3$  while  $[ABC] = [BC][AB]/2$  and  $[ABD] = [BD][AB]/2$ , so our equation's right-hand side is  $\frac{2[BCD]}{[BC][BD]} = \sin \angle CBD = \sin \theta$ .

**Lemma 7.** *Each of the derivatives  $\frac{\partial \theta}{\partial AB}, \frac{\partial \theta}{\partial AC}, \frac{\partial \theta}{\partial AD}$  is a degree-10 polynomial in the side lengths of  $ABCD$ , divided by  $[ABCD][ABC]^2[ABD]^2$ .*

*Proof.* Write  $[ABC]^2, [ABD]^2$  as polynomials in the side lengths using Heron's formula. Write  $[ABCD]^2$  as a polynomial in the side lengths as follows. We have  $36[ABCD]^2 = \det([\mathbf{AB}, \mathbf{AC}, \mathbf{AD}])^2 = \det(M)$  where  $M = [\mathbf{AB}, \mathbf{AC}, \mathbf{AD}]^T [\mathbf{AB}, \mathbf{AC}, \mathbf{AD}]$ . The entries of  $M$  are of the form  $\mathbf{u} \cdot \mathbf{v} = \frac{1}{2}(|\mathbf{u}|^2 + |\mathbf{v}|^2 - |\mathbf{u} - \mathbf{v}|^2)$ , which are polynomials in the side lengths. With Lemma 6, this gives  $\sin^2 \theta$  as a rational function of the side lengths.

Now  $\frac{\partial \theta}{\partial x} = \frac{\partial \sin \theta}{\partial x} / \sqrt{1 - \sin^2 \theta}$  for any variable  $x$ , so the square of this first derivative is a rational function. Computing it in SAGE [11] or another computer algebra system finds that for each  $x \in \{AB, AC, AD\}$ , this squared deriva-

tive has numerator the square of a degree-10 polynomial with denominator  $[ABCD]^2[ABC]^4[ABD]^4$ . The lemma is proved.

**Lemma 8.** *Each of the six second partial derivatives of  $\theta$  in  $AB, AC, AD$  is a degree-23 polynomial in the side lengths of  $ABCD$ , divided by  $[ABCD]^3[ABC]^4[ABD]^4$ .*

*Proof.* By Lemma 7, each first partial derivative is a degree-10 polynomial divided by  $[ABCD][ABC]^2[ABD]^2$ . Since  $[ABCD]^2, [ABC]^2, [ABD]^2$  are polynomials of degree 6, 4, 4 respectively, their logarithmic derivatives have themselves in the denominator and polynomials of degree 5, 3, 3 respectively in the numerator. The second partial derivatives therefore have an additional factor of  $[ABCD]^2[ABC]^2[ABD]^2$  in the denominator and an additional degree of 13 in the numerator, proving the lemma.

#### A.4 Proofs for Subsection 6.1

In this subsection we prove Lemma 9. The proof requires a lemma from singular spherical geometry.

**Lemma 15.** *Let  $C$  be a convex singular spherical  $n$ -gon with one interior vertex  $v$  of defect  $\kappa$  and each boundary vertex  $v_i$  a distance  $\alpha \leq vv_i \leq \beta \leq \pi/2$  from  $v$ . Then the perimeter  $\text{per}(C)$  is bounded by*

$$2\pi - \kappa - 2n(\pi/2 - \alpha) \leq \text{per}(C) \leq (2\pi - \kappa) \sin \beta.$$

*Proof.* Embed  $C$  in the singular spherical polygon  $B$  that results from removing a wedge of angle  $\kappa$  from a hemisphere.

To derive the lower bound, let the nearest point on the equator to each  $v_i$  be  $u_i$ , so that  $u_i v_i \leq \pi/2 - \alpha$ . Then by the triangle inequality,

$$\text{per}(C) = \sum_{ij} v_i v_j \geq \sum_{ij} u_i u_j - v_i u_i - u_j v_j \geq 2\pi - \kappa - 2n(\pi/2 - \alpha).$$

For the upper bound, let  $D$  be the singular spherical surface obtained as the  $\beta$ -disk about  $v$  in  $B$ . Then  $C$  can be obtained by cutting  $D$  in turn along the geodesic extension of each of the sides of  $C$ . Each of these cuts, because it is a geodesic, is the shortest path with its winding number and is therefore shorter than the boundary it replaces, so the perimeter only decreases in this process. Therefore  $\text{per}(C) \leq \text{per}(D) = (2\pi - \kappa) \sin \beta$ .

**Lemma 9.** *Given a polyhedral metric space  $M$ , there exists a radius assignment  $r$  with curvature skew  $\varepsilon_7 < \varepsilon_8/4\pi$ , maximum radius  $R = O(nD/\varepsilon_1\varepsilon_8)$ , and minimum defect-curvature gap  $\varepsilon_2 = \Omega(\varepsilon_1^2\varepsilon_8^3/n^2S^2)$ .*

*Proof.* Let  $r$  have the same value  $R$  on all vertices. We show that for sufficiently large  $R = O(nD/\varepsilon_1\varepsilon_8)$  the assignment  $r$  is valid and satisfies the required bounds on  $\varepsilon_2$  and  $\varepsilon_7$ . To do this it suffices to show that  $\varepsilon_2 \leq \delta_i - \kappa_i \leq \varepsilon_7\varepsilon_1$  for the desired  $\varepsilon_2, \varepsilon_7$  and each  $i$ .

For each vertex  $v_i$ , consider the singular spherical polygon  $C$  formed at  $v_i$  by the neighboring tetrahedra  $v_iOv_jv_k$ . Polygon  $C$  has one interior vertex at  $v_iO$  with defect  $\kappa_i$ , its perimeter is  $\sum_{jk} \angle v_jv_iv_k = 2\pi - \delta_i$ , and each vertex  $v_iv_k$  is convex. The spherical distance from the center  $v_iO$  to each vertex  $v_iv_k$  is  $\angle Ov_iv_k = \pi/2 - \Theta(v_iv_k/R)$ , which is at least  $\rho_{\min} \triangleq \pi/2 - \Theta(D/R)$  and at most  $\rho_{\max} \triangleq \pi/2 - \Theta(\ell/R)$ . Now by Lemma 15 above, we have

$$2\pi - \kappa_i - 2n(\pi/2 - \rho_{\min}) \leq 2\pi - \delta_i \leq (2\pi - \kappa_i) \sin \rho_{\max}.$$

The left-hand inequality implies

$$\delta_i - \kappa_i \leq 2n(\pi/2 - \rho_{\min}) = O(nD/R)$$

so that  $\delta_i - \kappa_i \leq (\varepsilon_8/4\pi)\varepsilon_1$  if  $R = \Omega(nD/\varepsilon_1\varepsilon_8)$  for a sufficiently large constant factor. The right-hand inequality then implies

$$\delta_i - \kappa_i \geq (2\pi - \delta_i) \frac{1 - \sin \rho_{\max}}{\sin \rho_{\max}} \geq \varepsilon_8(1 - \sin \rho_{\max}) = \Omega(\varepsilon_8\ell^2/R^2) = \Omega(\varepsilon_1^2\varepsilon_8^3/n^2S^2)$$

so that the  $\varepsilon_2$  bound holds.

## A.5 Proofs for Subsection 6.2

In this subsection we prove Lemmas 10 and 11.

**Lemma 10.**  $\varepsilon_3 > \ell d_1/R^2$ .

*Proof.*  $\varepsilon_3$  is the smallest angle  $\phi_{ij}$  from the apex  $O$  between any two vertices  $v_iv_j$ . Now  $v_iv_j \geq \ell$ , and the altitude from  $O$  to  $v_iv_j$  is at least  $d_1$ . Therefore  $\frac{1}{2}\ell d_1 \leq [Ov_iv_j] \leq \frac{1}{2} \sin \phi_{ij} R^2$ , so  $\phi_{ij} > \sin \phi_{ij} \geq \ell d_1/R^2$ .

**Lemma 11.**  $\varepsilon_5 > \varepsilon_2/6S$ .

*Proof.* Suppose that a surface triangle has an angle of  $\epsilon$ ; we want to show  $\epsilon > \varepsilon_2/6S$ . Let the largest angle of that triangle be  $\pi - \epsilon'$ . By the law of sines,  $\frac{\sin \epsilon'}{\sin \epsilon} \leq S$ , so  $\epsilon > \sin \epsilon \geq \sin \epsilon'/S > \epsilon'/3S$  since  $\epsilon' \leq 2\pi/3$  implies  $\sin \epsilon'/\epsilon' > 1/3$ . It therefore suffices to show that  $\epsilon' \geq \varepsilon_2/2$ .

Let the angle of size  $\pi - \epsilon'$  be at vertex  $i$ . Embed all of the tetrahedrons around  $Ov_i$  in space so that all the faces line up except for the one corresponding to

an edge  $e$  adjacent to this angle of  $\pi - \epsilon'$ . The two copies of  $e$  are separated by an angle of  $\kappa_i$ . Letting  $f$  be the other side forming this large angle, the angle between one copy of  $e$  and the copy of  $f$  is  $\pi - \epsilon'$ . Now the sum of all the angles around  $v_i$  is  $2\pi - \delta_i$ , so apply the triangle inequality for angles twice to deduce

$$\begin{aligned}
\epsilon_2 &\leq 2\pi - (2\pi - \delta_i) - \kappa_i \\
&\leq 2\pi - ((\pi - \epsilon') + \angle fe') - \kappa_i \\
&= \pi + \epsilon' - \angle fe' - \kappa_i \\
&\leq \pi + \epsilon' - ((\pi - \epsilon') - \kappa_i) - \kappa_i \\
&= 2\epsilon'.
\end{aligned}$$

### A.6 Proofs for Subsection 6.3

In this subsection we prove Lemmas 12, 13, and 14. The proofs require two additional lemmas about singular spherical polygons and matrices.

**Lemma 16.** *Let a convex singular spherical polygon have all exterior angles at least  $\gamma$  and all side lengths between  $c$  and  $2\pi - c$ . Then its perimeter is at most  $2\pi - \Omega(\gamma^2 c)$ .*

*Proof.* This is an effective version of Lemma 5.4 on pages 45–46 of [2], and we follow their proof. The proof in [2] shows that the perimeter is in general bounded by the perimeter in the nonsingular case. In this case consider any edge  $AB$  of the polygon, and observe that since the polygon is contained in the triangle  $ABC$  with exterior angles  $\gamma$  at  $A, B$  its perimeter is bounded by this triangle's perimeter. Since  $c \leq AB \leq 2\pi - c$ , the bound follows by straightforward spherical geometry.

**Lemma 17.** *Let  $S$  be a singular spherical metric with vertices  $\{v_i\}_i$ , and let  $C$  be the singular spherical polygon consisting of the triangles about some distinguished vertex  $v_0$ . Suppose  $C$  has  $k$  convex vertices, each with an interior angle at least  $\pi - \epsilon$  for some  $\epsilon > 0$  and an exterior angle no more than  $\pi$ . Then*

$$\kappa_0 + 2\epsilon k \geq \left(1 - \frac{\text{per}(C)}{2\pi}\right) \sum_{i \neq 0} \kappa_i.$$

*Proof.* We reduce to Lemma 5.5 from [2] by induction. If  $k = 0$ , so that all vertices of  $C$  have interior angle at least  $\pi$ , then our statement is precisely theirs.

Otherwise, let  $v_i$  be a vertex of  $C$  with interior angle  $\pi - \theta \in [\pi - \epsilon, \pi)$ . Draw the geodesic from  $v_i$  to  $v_0$ , and insert along this geodesic a pair of spherical

triangles each with angle  $\theta/2$  at  $v_i$  and angle  $\kappa_0/2$  at  $v_0$ , meeting at a common vertex  $v'_0$ . The polygon  $C'$  and triangulation  $S'$  that result from adding these two triangles satisfy all the same conditions but with  $k - 1$  convex vertices on  $C'$ , so

$$\kappa'_0 + 2\varepsilon(k - 1) \geq \left(1 - \frac{\text{per}(C')}{2\pi}\right) \sum_{i \neq 0} \kappa'_i.$$

Now  $C'$  and  $C$  have the same perimeter,  $\kappa'_0 \leq \kappa_0 + \theta \leq \kappa_0 + \varepsilon$ ,  $\kappa'_i = \kappa_i - \theta \geq \kappa_i - \varepsilon$ , and  $\kappa'_j = \kappa_j$  for  $j \notin \{0, i\}$ , so it follows that

$$\kappa_0 + 2\varepsilon k \geq \kappa'_0 + (2k - 1)\varepsilon \geq \left(1 - \frac{\text{per}(C')}{2\pi}\right) \sum_{i \neq 0} \kappa_i$$

as claimed.

**Lemma 12.**  $d_2 = \Omega(D\varepsilon_1\varepsilon_4\varepsilon_5^2\varepsilon_8/(nS^4))$ .

*Proof.* This is an effective version of Lemma 4.8 of [2], on whose proof this one is based.

Let  $i = \arg \min_i Ov_i$ , so that  $Ov_i = d_2$ , and suppose that  $d_2 = O(D\varepsilon_1\varepsilon_4\varepsilon_5^2\varepsilon_8/(nS^4))$  with a small constant factor. We consider the singular spherical polygon  $C$  formed at the apex  $O$  by the tetrahedra about  $Ov_i$ . First we show that  $C$  is concave or nearly concave at each of its vertices, so that it satisfies the hypothesis of Lemma 17. Then we apply Lemma 17 and use the fact that the ratios of the  $\kappa_j$  are within  $\varepsilon_7 \leq \varepsilon_8/4\pi$  of those of the  $\delta_j$  to get a contradiction.

Consider a vertex of  $C$ , the ray  $Ov_j$ . Let  $v_iv_jv_k, v_jv_iv_l$  be the triangles in  $T$  adjacent to  $v_iv_j$ , and embed the two tetrahedra  $Ov_iv_jv_k, Ov_jv_iv_l$  in  $\mathbb{R}^3$ . The angle of  $C$  at  $Ov_j$  is the dihedral angle  $v_kOv_jv_l$ .

By convexity, the dihedral angle  $v_kv_iv_jv_l$  contains  $O$ , so if  $O$  is on the same side of plane  $v_kv_jv_l$  as  $v_i$  is then the dihedral angle  $v_kOv_jv_l$  does not contain  $v_i$  and is a reflex angle for  $C$ . Otherwise, the distance from  $O$  to this plane is at most  $Ov_i = d_2$ , and we will bound the magnitude of  $\angle v_kOv_jv_l$ .

By Lemma 6,

$$\sin \angle v_kOv_jv_l = \frac{3 [Ov_kv_jv_l][Ov_j]}{2 [Ov_jv_k][Ov_jv_l]}.$$

Now  $[Ov_kv_jv_l] \leq d_2[v_kv_jv_l]/3 = O(d_2D^2)$  and  $[Ov_j] \leq [Ov_i] + [v_iv_j] \leq D + d_2$ . On the other hand  $[Ov_jv_k] = (1/2)[Ov_j][Ov_k] \sin \angle v_jOv_k$ , and  $[Ov_j], [Ov_k] \geq \ell - d_2$  while  $\angle v_kv_iv_j \leq \angle v_kv_kO + \angle v_kOv_j + \angle Ov_jv_i \leq \angle v_kOv_j + O(d_2/D)$  so that  $\angle v_jOv_k \geq \varepsilon_5 - O(d_2/D)$ , so  $[Ov_jv_k] = \Omega(\ell^2\varepsilon_5)$ , and similarly  $[Ov_jv_l]$ . Therefore  $\sin \angle v_kOv_jv_l = O(d_2D^3/(\ell^4\varepsilon_5^2)) = O(\varepsilon_1\varepsilon_4\varepsilon_8/n)$ , and the angle of  $C$  at  $Ov_i$  is

$$\angle v_kOv_jv_l = O(\varepsilon_1\varepsilon_4\varepsilon_8/n).$$

On the other hand observe that  $\text{per}(C) = \sum_{jk, v_i v_j v_k \in F(T)} \angle v_j O v_k \leq \sum_{jk} (\angle v_j v_i v_k + O(d_2/D)) = 2\pi - \delta_i + O(nd_2/D)$ .

Now apply Lemma 17 to deduce that

$$\kappa_i + O(\varepsilon_1 \varepsilon_4 \varepsilon_8) \geq \left(1 - \frac{\text{per}(C)}{2\pi}\right) \sum_{j \neq i} \kappa_j \geq \left(\frac{\delta_i}{2\pi} - O(nd_2/D)\right) \sum_{j \neq i} \kappa_j$$

so that

$$\begin{aligned} \frac{\kappa_i}{\delta_i} + O(\varepsilon_4 \varepsilon_8) &\geq \left(\frac{1}{2\pi} - O(nd_2/\varepsilon_1 D)\right) \sum_{j \neq i} \kappa_j \\ &\geq (1 + o(\varepsilon_8)) \frac{1}{2\pi} \left(\min_j \frac{\kappa_j}{\delta_j}\right) \sum_{j \neq i} \delta_j \\ &= (1 + o(\varepsilon_8)) \frac{4\pi - \delta_i}{2\pi} \left(\min_j \frac{\kappa_j}{\delta_j}\right) \\ &\geq (1 + o(\varepsilon_8))(1 + \varepsilon_8/2\pi) \left(\min_j \frac{\kappa_j}{\delta_j}\right) \end{aligned}$$

so that since  $\kappa_i/\delta_i = \Omega(\varepsilon_4)$ ,

$$\frac{\kappa_i}{\delta_i} \left(\min_j \frac{\kappa_j}{\delta_j}\right)^{-1} \geq (1 + O(\varepsilon_8))^{-1} (1 + \varepsilon_8/2\pi)$$

which for a small enough constant factor on  $d_2$  and hence on the  $O(\varepsilon_8)$  term makes  $\varepsilon_7 > \varepsilon_8/(4\pi)$ , which is a contradiction.

**Lemma 13.**  $d_1 = \Omega(\varepsilon_4^2 \varepsilon_5^2 d_2^2 / DS^2) = \Omega(D \varepsilon_1^2 \varepsilon_4^4 \varepsilon_5^6 \varepsilon_8^2 / (n^2 S^{10}))$ .

*Proof.* This is an effective version of Lemma 4.6 of [2], on whose proof this one is based.

Let  $O$  be distance  $d_1$  from edge  $v_i v_j$ , which neighbors faces  $v_i v_j v_k, v_j v_i v_l \in F(T)$ . Consider the spherical quadrilateral  $D$  formed at  $O$  by the two tetrahedra  $Ov_i v_j v_k, Ov_j v_i v_l$ , and the singular spherical quadrilateral  $C$  formed by all the other tetrahedra. We will show the perimeter of  $C$  is nearly  $2\pi$  for small  $d_1$  and apply Lemma 16 to deduce a bound. This requires also upper and lower bounds on the side lengths of  $C$  and a lower bound on its exterior angles.

In triangle  $Ov_i v_j$ , let the altitude from  $O$  have foot  $q$ ; then  $Oq = d_1$  while  $v_i O, v_j O \geq d_2$ , so  $\angle v_j v_i q, \angle v_i v_j q = O(d_1/d_2)$ . Also,  $qv_i, qv_j \geq d_2 - d_1$ , so  $q$  is at least distance  $(d_2 - d_1) \sin \varepsilon_5$  from any of  $v_i v_k, v_k v_j, v_j v_l, v_l v_i$ , and  $O$  is at least  $(d_2 - d_1) \sin \varepsilon_5 - d_1 = \Omega(d_2 \varepsilon_5)$  from each of these sides.

Now  $\angle v_i O v_j = \pi - O(d_1/d_2)$  is the distance on the sphere between opposite vertices  $Ov_i, Ov_j$  of  $D$ , so by the triangle inequality the perimeter of  $D$  is at least  $2\pi - O(d_1/d_2)$ . Each side of  $C$  is at least  $\Omega(\varepsilon_5)$  and at most  $\pi - \Omega(\varepsilon_5 d_2/D)$ .

In spherical quadrilateral  $D$ , the two opposite angles  $\angle v_k O v_i v_l$ ,  $\angle v_l O v_j v_k$  are each within  $O(d_1/\varepsilon_5 d_2)$  of the convex  $\angle v_k v_i v_j v_l$  and therefore either reflex for  $D$  or else at least  $\pi - O(d_1/\varepsilon_5 d_2)$ . To bound the other two angles  $\angle v_i O v_l v_j$ ,  $\angle v_j O v_k v_i$ , let the smaller of these be  $\theta$ ; then by Lemma 6,

$$\pi - \theta = O(\sin \theta) = O\left(\frac{(D^2 d_1) D}{(\varepsilon_5 d_2 D/S)^2}\right) = O\left(\frac{S^2 D d_1}{\varepsilon_5^2 d_2^2}\right).$$

Now there are two cases. In one case,  $d_1 = \Omega(\varepsilon_4 \varepsilon_5^2 d_2^2 / DS^2)$ . In the alternative, we find that each angle of  $D$  is at least  $\pi - \varepsilon_4/2$  and each angle of  $C$  at most  $\pi - \varepsilon_4/2$ . In the latter case applying Lemma 16 to  $C$  finds that  $2\pi - O(d_1/d_2) = 2\pi - \Omega(\varepsilon_4^2 \varepsilon_5 d_2 / D)$  so that  $d_1 = \Omega(\varepsilon_4^2 \varepsilon_5 d_2^2 / D)$ .

In either case  $d_1 = \Omega(\min(\varepsilon_4 \varepsilon_5^2 d_2^2 / DS^2, \varepsilon_4^2 \varepsilon_5 d_2^2 / D)) = \Omega(\varepsilon_4^2 \varepsilon_5^2 d_2^2 / DS^2)$ , and the bound on  $d_2$  from Lemma 12 finishes the proof.

**Lemma 14.**  $d_0 = \Omega(D \varepsilon_1^4 \varepsilon_4^9 \varepsilon_5^{12} \varepsilon_8^4 / (n^4 S^{22}))$ .

*Proof.* This is an effective version of Lemma 4.5 of [2], on whose proof this one is based.

Let  $O$  be distance  $d_0$  from triangle  $v_i v_j v_k \in F(T)$ . Consider the singular spherical polygon  $C$  cut out at  $O$  by all the tetrahedra other than  $O v_i v_j v_k$ . We show lower and upper bounds on the side lengths of  $C$  and lower bounds on its exterior angles, show the perimeter  $\text{per}(C)$  is near  $2\pi$  for small  $d_0$ , and apply Lemma 16 to derive a bound.

The perimeter of  $C$  is the total angle about  $O$  on the faces of the tetrahedron  $O v_i v_j v_k$ , which is  $2\pi - O(d_0^2/d_1^2)$ . Each side of  $C$  is at least  $\Omega(\varepsilon_5)$  and at most  $\pi - \Omega(d_1/D)$ .

Let  $\theta$  be the smallest dihedral angle of  $\angle v_i O v_j v_k$ ,  $\angle v_j O v_k v_i$ ,  $\angle v_k O v_i v_j$ . Then by Lemma 6,

$$\pi - \theta = O(\sin \theta) = O\left(\frac{(D^2 d_0) D}{(d_1 D/S)^2}\right) = O\left(\frac{S^2 D d_0}{d_1^2}\right).$$

Now there are two cases. If  $\theta \leq \pi - \varepsilon_4/2$ , then it follows immediately that  $d_0 = \Omega(d_1^2 \varepsilon_4 / (S^2 D))$ . Otherwise,  $\theta > \pi - \varepsilon_4/2$ , so the interior angles of  $C$  are more than  $\varepsilon_4/2$ . Applying Lemma 16, the perimeter  $\text{per}(C)$  is at most  $2\pi - \Omega(\min(\varepsilon_4^2 \varepsilon_5, \varepsilon_4^2 d_1 / D))$ , so that  $d_0 = \Omega(\min(d_1 \varepsilon_4 \varepsilon_5^{1/2}, d_1^{3/2} \varepsilon_4 D^{-1/2}))$ . The bound on  $d_1$  from Lemma 13 finishes the proof.