

# Logic Synthesis and Verification

Jie-Hong Roland Jiang  
江介宏

Department of Electrical Engineering  
National Taiwan University



Fall 2012

1

## Two-Level Logic Minimization (1/2)

Reading:

*Logic Synthesis in a Nutshell*

Section 3 (§3.1-§3.2)

most of the following slides are by  
courtesy of Andreas Kuehlmann

2

# Quine-McCluskey Procedure


- Given  $G$  and  $D$  (covers for  $\mathfrak{S} = (f, d, r)$  and  $d$ , respectively), find a minimum cover  $G^*$  of primes where:
 
$$f \subseteq G^* \subseteq f+d \quad (G^* \text{ is a prime cover of } \mathfrak{S})$$
- Q-M Procedure:
  1. Generate all primes of  $\mathfrak{S}$ ,  $\{P_j\}$  (i.e. primes of  $(f+d) = G+D$ )
  2. Generate all minterms  $\{m_i\}$  of  $f = G \wedge \neg D$
  3. Build Boolean matrix  $B$  where
 
$$B_{ij} = 1 \text{ if } m_i \in P_j$$

$$= 0 \text{ otherwise}$$
  4. Solve the minimum column covering problem for  $B$  (unate covering problem)

3

# Complexity

- $\sim 2^n$  minterms;  $\sim 3^n/n$  primes




- There are  $O(2^n)$  rows and  $\Omega(3^n/n)$  columns. Moreover, minimum covering problem is NP-complete. (Hence the complexity can probably be double exponential in size of input, i.e. difficulty is  $O(2^{3^n})$ )

4

# Two-Level Logic Minimization

## Example



$$F = \bar{x} \bar{y} z w + \bar{x} y \bar{z} w + x \bar{y} z w + x y z w \quad (\text{cover of } \mathfrak{F})$$

$$D = \bar{y} z + x y w + \bar{x} y \bar{z} w + x \bar{y} w + \bar{x} y \bar{z} w \quad (\text{cover of } d)$$

Primes:  $\bar{y} + w + \bar{x} \bar{z}$

Covering Table

Solution:  $\{1,2\} \Rightarrow \bar{y} + w$  is a minimum prime cover (also  $w + \bar{x} \bar{z}$ )

5

# Covering Table

	$\bar{y}$	$w$	$\bar{x} \bar{z}$	Primes of f+d
$\bar{x} \bar{y} \bar{z} \bar{w}$	1	0	1	
$\bar{x} y \bar{z} w$	0	1	1	
$x \bar{y} \bar{z} w$	1	1	0	
$\bar{x} y z w$	0	1	0	Row singleton (essential minterm)

↑  
Essential prime

Definition. An essential prime is a prime that covers an onset minterm of f not covered by any other primes.

6

# Covering Table

## Row Equality

### □ Row equality:

- In practice, many rows in a covering table are identical. That is, there exist minterms that are contained in the same set of primes.

### ■ Example

$m_1$	0101101
$m_2$	0101101

7

# Covering Table

## Row and Column Dominance

### □ Row dominance:

- A row  $i_1$  whose set of primes is contained in the set of primes of row  $i_2$  is said to **dominate**  $i_2$ .

### ■ Example

$i_1$	011010
$i_2$	011110

- $i_1$  dominates  $i_2$
- Can remove row  $i_2$  because have to choose a prime to cover  $i_1$ , and any such prime also covers  $i_2$ . So  $i_2$  is automatically covered.

8

# Covering Table

## Row and Column Dominance

### □ Column dominance:

- A *column*  $j_1$  whose rows are a superset of another *column*  $j_2$  is said to **dominate**  $j_2$ .

### ■ Example

$j_1$	$j_2$
1	0
0	0
1	1
0	0
1	1

- $j_1$  dominates  $j_2$
- We can remove column  $j_2$  since  $j_1$  covers all those rows and more. We would never choose  $j_2$  in a minimum cover since it can always be replaced by  $j_1$ .

9

# Covering Table

## Table Reduction


1. Remove all rows covered by essential primes (columns in row singletons). Put these primes in the cover  $G$ .
2. Group identical rows together and remove dominated rows.
3. Remove dominated columns. For equal columns, keep one prime to represent them.
4. Newly formed row singletons define **induced essential primes**.
5. Go to 1 if covering table decreased.

- The resulting reduced covering table is called the **cyclic core**. This has to be solved (**unate covering problem**). A minimum solution is added to  $G$ . The resulting  $G$  is a minimum cover.

10

# Covering Table Table Reduction

## Example



11

# Solving Cyclic Core


Best known method (for unate covering) is **branch and bound** with some clever bounding heuristics

## Independent Set Heuristic:

Find a maximum set  $I$  of "independent" rows. Two rows  $B_{i_1}, B_{i_2}$  are independent if **not**  $\exists j$  such that  $B_{i_1j} = B_{i_2j} = 1$ . (They have **no column in common.**)

### Example

A covering matrix  $B$  rearranged with independent sets first



12

# Solving Cyclic Core

□ Lemma:

$$|\text{Solution of Covering}| \geq |\mathcal{F}|$$

$m_1$  must be covered by one of the three columns

$m_1$	1 1 1	0
$m_2$	1 1 1 1	0
$m_3$	1 1	0
	A	C

13

# Solving Cyclic Core

□ Heuristic algorithm:

- Let  $\mathcal{F} = \{I_1, I_2, \dots, I_k\}$  be the independent set of rows
- 1. choose  $j \in I_i$  such that column  $j$  covers the most rows of A. Put  $P_j$  in  $G$
- 2. eliminate all rows covered by column  $j$
- 3.  $\mathcal{F} \leftarrow \mathcal{F} \setminus \{I_i\}$
- 4. go to 1 if  $|\mathcal{F}| > 0$
- 5. If B is empty, then done (in this case achieve minimum solution because of the lower bound of previous lemma attained - **IMPORTANT**)
- 6. If B is not empty, choose an independent set of B and go to 1

1 1 1	0
1 1 1 1	0
1 1	0
A	C

14

# Prime Generation for Single-Output Function

## Tabular method

(based on *consensus* operation, or  $\nabla$ ):

- Start with minterm canonical form of  $F$
- Group *pairs* of adjacent minterms into cubes
- Repeat merging cubes until no more merging possible; mark ( $\checkmark$ ) + remove all covered cubes.
- Result: set of *primes* of  $f$ .

## Example

$$F = x' y' + w x y + x' y z' + w y' z$$

$$F = x' y' + w x y + x' y z' + w y' z$$

$w' x' y' z' \checkmark$	$w' x' y' \checkmark$	$x' y' \checkmark$
	$w' x' z' \checkmark$	$x' z' \checkmark$
	$x' y' z' \checkmark$	
$w' x' y' z \checkmark$	$x' y' z \checkmark$	
$w' x' y z' \checkmark$	$x' y z' \checkmark$	
$w x' y' z' \checkmark$	$w x' y' \checkmark$	
	$w x' z' \checkmark$	
$w x' y' z \checkmark$	$w y' z \checkmark$	
$w x' y z' \checkmark$	$w y z' \checkmark$	
$w x y z' \checkmark$	$w x y \checkmark$	
$w x y' z \checkmark$	$w x z \checkmark$	
$w x y z \checkmark$		


Courtesy: Maciej Ciesielski, UMASS

15

# Prime Generation for Multi-Output Function

- Similar to *single-output* function, except that we should include also the **primes of the products of individual functions**

## Example



$x y z$	$f_1 f_2$
0-0	01
011	11
1-1	10

$x y z$	$f_1 f_2$
0-0	01
01-	01
-11	10
1-1	10

Can also represent it as:


16



# Prime Generation

## Example

- Modification from single-output case: When two adjacent implicants are merged, the output parts are **intersected**



There are five primes for this two-output function  
- What is the min cover ?

17

# Minimize Multi-Output Cover


## Example

- List multiple-output primes
- Create a covering table
- Solve

$$\begin{aligned}
 p_1 &= 0 1 1 | 11 \\
 p_2 &= 0 - 0 | 01 \\
 p_3 &= 0 1 - | 01 \\
 p_4 &= - 1 1 | 10 \\
 p_5 &= 1 - 1 | 10
 \end{aligned}$$

Min cover has 3 primes:

$$F = \{ p_1, p_2, p_5 \}$$



18

# Prime Generation Usingunate Recursive Paradigm

- Apply **unate recursive paradigm** with the following **merge step**
  - (Assume we have just generated all primes of  $f_{x_i}$  and  $f_{\neg x_i}$ )

- **Theorem.**

$p$  is a prime of  $f$  iff  $p$  is **maximal** (in terms of containment) among the set consisting of

- $P = x_i q$ ,  $q$  is a prime of  $f_{x_i}$ ,  $q \not\subseteq f_{\neg x_i}$
- $P = x_i' r$ ,  $r$  is a prime of  $f_{\neg x_i}$ ,  $r \not\subseteq f_{x_i}$
- $P = q r$ ,  $q$  is a prime of  $f_{x_i}$ ,  $r$  is a prime of  $f_{\neg x_i}$

19

# Prime Generation Usingunate Recursive Paradigm

- **Example**

- Assume  $q = abc$  is a prime of  $f_{x_i}$ . Form  $p = x_i abc$ .
- Suppose  $r = ab$  is a prime of  $f_{\neg x_i}$ . Then  $x_i' ab$  is an implicant of  $f$ .

$$f = x_i abc + x_i' ab + abc + \dots$$

- Thus  $abc$  and  $x_i' ab$  are implicants, so  $x_i abc$  is not prime.
- **Note:**  $abc$  is prime because if not,  $ab \subseteq f$  (or  $ac$ , or  $bc$ ) contradicting  $abc$  prime of  $f_{x_i}$ .
- **Note:**  $x_i' ab$  is prime, since if not then either  $ab \subseteq f$ ,  $x_i' a \subseteq f$ ,  $x_i' b \subseteq f$ . The first contradicts  $abc$  prime of  $f_{x_i}$  and the second and third contradict  $ab$  prime of  $f_{\neg x_i}$ .

20

# Summary

## □ Quine-McCluskey Method:

1. Generate cover of all primes  $G = p_1 + p_2 + \dots + p_{3^n/n}$
2. Make G irredundant (in optimum way)

- Q-M is **exact**, i.e., it gives an exact minimum

## □ Heuristic Methods:

1. Generate (somehow) a cover of  $\mathfrak{F}$  using some of the primes  $G = p_{i_1} + p_{i_2} + \dots + p_{i_k}$
2. Make G irredundant (maybe not optimally)
3. Keep best result - try again (i.e. go to 1)