CSE 203B Week 5 Discuss Session

Xinyuan Wang

02/06/2019

Contents

- Convex Optimization Problems (Ref. Chap 4)
  - Optimization problem in standard form
  - Convex optimization problem
  - Linear optimization
  - Quadratic optimization
  - Geometric programming

Optimization problem in standard form

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \ldots, m \\ & h_i(x) = 0, \quad i = 1, \ldots, p \end{array}$$

- $x \in \mathbf{R}^n$ is the optimization variable
- $f_0 : \mathbf{R}^n \to \mathbf{R}$ is the objective or cost function
- $f_i : \mathbf{R}^n \to \mathbf{R}$, $i = 1, \ldots, m$, are the inequality constraint functions
- $h_i : \mathbf{R}^n \to \mathbf{R}$ are the equality constraint functions

optimal value:

$$p^\star = \inf\{f_0(x) \mid f_i(x) \leq 0,\ i = 1, \ldots, m,\ h_i(x) = 0,\ i = 1, \ldots, p\}$$

- $p^\star = \infty$ if problem is infeasible (no x satisfies the constraints)
- $p^\star = -\infty$ if problem is unbounded below

Optimal and locally optimal points

x is **feasible** if $x \in \mathbf{dom}\, f_0$ and it satisfies the constraints

a feasible x is **optimal** if $f_0(x) = p^\star$; $X_{\rm opt}$ is the set of optimal points

x is **locally optimal** if there is an $R > 0$ such that x is optimal for

$$\begin{array}{ll} \text{minimize (over } z) & f_0(z) \\ \text{subject to} & f_i(z) \leq 0, \quad i = 1, \ldots, m, \quad h_i(z) = 0, \quad i = 1, \ldots, p \\ & \|z - x\|_2 \leq R \end{array}$$

examples (with $n = 1$, $m = p = 0$)

- $f_0(x) = 1/x$, $\mathbf{dom}\, f_0 = \mathbf{R}_{++}$: $p^\star = 0$, **opt. value not achieved**
- $f_0(x) = -\log x$, $\mathbf{dom}\, f_0 = \mathbf{R}_{++}$: $p^\star = -\infty$ **unbounded below**
- $f_0(x) = x \log x$, $\mathbf{dom}\, f_0 = \mathbf{R}_{++}$: $p^\star = -1/e$, $x = 1/e$ is optimal **unique opt.**
- $f_0(x) = x^3 - 3x$, $p^\star = -\infty$, local optimum at $x = 1$ **local opt. ≠ global op.**

Implicit constraints

the standard form optimization problem has an **implicit constraint**

$$x \in \mathcal{D} = \bigcap_{i=0}^{m} \mathbf{dom}\, f_i \cap \bigcap_{i=1}^{p} \mathbf{dom}\, h_i,$$

- we call $\mathcal{D}$ the **domain** of the problem
- the constraints $f_i(x) \leq 0$, $h_i(x) = 0$ are the explicit constraints
- a problem is **unconstrained** if it has no explicit constraints $(m = p = 0)$

example:

$$\text{minimize} \quad f_0(x) = -\sum_{i=1}^{k} \log(b_i - a_i^T x)$$

is an unconstrained problem with implicit constraints $a_i^T x < b_i$

Feasibility problem

$$\begin{array}{ll} \text{find} & x \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \ldots, m \\ & h_i(x) = 0, \quad i = 1, \ldots, p \end{array}$$

can be considered a special case of the general problem with $f_0(x) = 0$:

$$\begin{array}{ll} \text{minimize} & 0 \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \ldots, m \\ & h_i(x) = 0, \quad i = 1, \ldots, p \end{array}$$

- $p^\star = 0$ if constraints are feasible; any feasible x is optimal
- $p^\star = \infty$ if constraints are infeasible

Convex optimization problem

standard form convex optimization problem

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \ldots, m \\ & a_i^T x = b_i, \quad i = 1, \ldots, p \end{array}$$

- $f_0, f_1, \ldots, f_m$ are convex; equality constraints are affine
- problem is *quasiconvex* if f_0 is quasiconvex (and $f_1, \ldots, f_m$ convex)

often written as

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \ldots, m \\ & Ax = b \end{array}$$

important property: feasible set of a convex optimization problem is convex

Local and global optima

★ any locally optimal point of a convex problem is (globally) optimal

proof: suppose x is locally optimal, but there exists a feasible y with $f_0(y) < f_0(x)$

x locally optimal means there is an $R > 0$ such that

$$z \text{ feasible}, \quad \|z - x\|_2 \leq R \quad \Longrightarrow \quad f_0(z) \geq f_0(x)$$

consider $z = \theta y + (1 - \theta)x$ with $\theta = R/(2\|y - x\|_2)$

- $\|y - x\|_2 > R$, so $0 < \theta < 1/2$
- z is a convex combination of two feasible points, hence also feasible
- $\|z - x\|_2 = R/2$ and

$$f_0(z) \leq \theta f_0(y) + (1 - \theta) f_0(x) < f_0(x)$$

which contradicts our assumption that x is locally optimal

Optimality criterion for differentiable f_0

★ x is optimal if and only if it is feasible and

$$\nabla f_0(x)^T(y-x) \geq 0 \quad \text{for all feasible } y$$

Proof: use first-order condition of the convex functions



if nonzero, $\nabla f_0(x)$ defines a supporting hyperplane to feasible set X at x

Optimality criterion for differentiable f_0

- **unconstrained problem**: x is optimal if and only if

$$x \in \mathbf{dom}\, f_0, \qquad \nabla f_0(x) = 0$$

Practice: prove the optimality criterion for convex opt. problems

- **equality constrained problem**

$$\text{minimize} \quad f_0(x) \quad \text{subject to} \quad Ax = b$$

x is optimal if and only if there exists a ν such that

$$x \in \mathbf{dom}\, f_0, \qquad Ax = b, \qquad \nabla f_0(x) + A^T\nu = 0$$

- **minimization over nonnegative orthant**

$$\text{minimize} \quad f_0(x) \quad \text{subject to} \quad x \succeq 0$$

x is optimal if and only if

$$x \in \mathbf{dom}\, f_0, \qquad x \succeq 0, \qquad \begin{cases} \nabla f_0(x)_i \geq 0 & x_i = 0 \\ \nabla f_0(x)_i = 0 & x_i > 0 \end{cases}$$

example

Example 4.5 *Unconstrained quadratic optimization.* Consider the problem of minimizing the quadratic function

$$f_0(x) = (1/2)x^T Px + q^T x + r,$$

where $P \in \mathbf{S}_+^n$ (which makes f_0 convex). The necessary and sufficient condition for x to be a minimizer of f_0 is

$$\nabla f_0(x) = Px + q = 0.$$

Several cases can occur, depending on whether this (linear) equation has no solutions, one solution, or many solutions.

- If $q \notin \mathcal{R}(P)$, then there is no solution. In this case f_0 is unbounded below.
- If $P \succ 0$ (which is the condition for f_0 to be strictly convex), then there is a unique minimizer, $x^\star = -P^{-1}q$.
- If P is singular, but $q \in \mathcal{R}(P)$, then the set of optimal points is the (affine) set $X_{\mathrm{opt}} = -P^\dagger q + \mathcal{N}(P)$, where $P^\dagger$ denotes the pseudo-inverse of P (see §A.5.4).

Equivalent convex problems

Two problems are ***equivalent*** if the solution of one is readily obtained from the solution of the other, and vice-versa

The transformations preserve **convexity** (Chap 4.2.4)

- Eliminating equality constraints
- Introducing equality constraints
- Introducing slack variables for linear inequalities
- Epigraph form
- Minimizing over some variables

Linear program (LP)

$$\begin{array}{ll} \text{minimize} & c^T x + d \\ \text{subject to} & Gx \preceq h \\ & Ax = b \end{array}$$

- convex problem with affine objective and constraint functions
- feasible set is a polyhedron



- **More examples in Chap 4.3**
- **Linear fractional program : transform to LP**

Quadratic program (QP)

$$\begin{array}{ll} \text{minimize} & (1/2)x^T Px + q^T x + r \\ \text{subject to} & Gx \preceq h \\ & Ax = b \end{array}$$

- $P \in \mathbf{S}_+^n$, so objective is convex quadratic
- minimize a convex quadratic function over a polyhedron



- **More examples in Chap 4.4**

Geometric programming

monomial function

$$f(x) = cx_1^{a_1}x_2^{a_2}\cdots x_n^{a_n}, \qquad \mathbf{dom}\, f = \mathbf{R}^n_{++}$$

with $c > 0$; exponent a_i can be any real number

posynomial function: sum of monomials

$$f(x) = \sum_{k=1}^{K} c_k x_1^{a_{1k}} x_2^{a_{2k}} \cdots x_n^{a_{nk}}, \qquad \mathbf{dom}\, f = \mathbf{R}^n_{++}$$

geometric program (GP)

$$\begin{array}{lll} \text{minimize} & f_0(x) & \\ \text{subject to} & f_i(x) \leq 1, & i = 1, \ldots, m \\ & h_i(x) = 1, & i = 1, \ldots, p \end{array}$$

with f_i posynomial, h_i monomial

Geometric program in convex form

change variables to $y_i = \log x_i$, and take logarithm of cost, constraints

- monomial $f(x) = cx_1^{a_1} \cdots x_n^{a_n}$ transforms to

$$\log f(e^{y_1}, \ldots, e^{y_n}) = a^T y + b \qquad (b = \log c)$$

- posynomial $f(x) = \sum_{k=1}^{K} c_k x_1^{a_{1k}} x_2^{a_{2k}} \cdots x_n^{a_{nk}}$ transforms to

$$\log f(e^{y_1}, \ldots, e^{y_n}) = \log \left(\sum_{k=1}^{K} e^{a_k^T y + b_k} \right) \qquad (b_k = \log c_k)$$

- geometric program transforms to convex problem

$$\begin{array}{ll} \text{minimize} & \log \left(\sum_{k=1}^{K} \exp(a_{0k}^T y + b_{0k}) \right) \\ \text{subject to} & \log \left(\sum_{k=1}^{K} \exp(a_{ik}^T y + b_{ik}) \right) \leq 0, \quad i = 1, \ldots, m \\ & Gy + d = 0 \end{array}$$

- **More examples in Chap 4.5**